DAMTP/02-114
hep-th/0209056

On Short and Semi-Short Representations for Four Dimensional Superconformal Symmetry

F.A. Dolan and H. Osborn†

Department of Applied Mathematics and Theoretical Physics,
Silver Street, Cambridge, CB3 9EW, England

arXiv:hep-th/0209056v2 7 Sep 2002

Possible short and semi-short representations for $\mathcal{N}=2$ and $\mathcal{N}=4$ superconformal symmetry in four dimensions are discussed. For $\mathcal{N}=4$ the well known short supermultiplets whose lowest dimension conformal primary operators correspond to $\frac{1}{2}$-BPS or $\frac{1}{4}$-BPS states and are scalar fields belonging to the $SU(4)_r$ symmetry representations $[0,p,0]$ and $[q,p,q]$ and having scale dimension $\Delta=p$ and $\Delta=2q+p$ respectively are recovered. The representation content of semi-short multiplets, which arise at the unitarity threshold for long multiplets, is discussed. It is shown how, at the unitarity threshold, a long multiplet can be decomposed into four semi-short multiplets. If the conformal primary state is spinless one of these becomes a short multiplet. For $\mathcal{N}=4$ a $\frac{1}{4}$-BPS multiplet need not have a protected dimension unless the primary state belongs to a $[1,p,1]$ representation.

PACS no: 11.25.Hf

Keywords: superconformal symmetry, supermultiplets, BPS.

† address for correspondence: Trinity College, Cambridge, CB2 1TQ, England
emails: fad20@damtp.cam.ac.uk and ho@damtp.cam.ac.uk

1. Introduction

Since the discovery of the AdS/CFT correspondence there has been a resurgence of interest in superconformal quantum field theories in four dimensions. The maximal $\mathcal{N} = 4$ theories for gauge group $SU(N)$ with coupling g have been of particular interest and, as is now well known, the supergravity approximation to type IIB string theories on $AdS_5 \times S^5$ gives direct information in the limit $N \to \infty$ for large λ where $\lambda = g^2 N$. The $\mathcal{N} = 4$ theory is superconformal for any g and it is possible to also obtain perturbative results valid for small λ. Of particular interest are correlation functions of chiral primary operators which may be written as $\mathcal{C}^I{}_{r_1 \dots r_p} \mathrm{tr}(X_{r_1} \dots X_{r_p})$ where $\mathcal{C}^I{}_{r_1 \dots r_p}$ are a set of symmetric traceless $SO(6)$ tensors and X_r are the lowest dimension scalar fields in the $\mathcal{N} = 4$ theory, belonging to the adjoint representation of the gauge group and the 6-dimensional representation of the R-symmetry group $SO(6)_R \simeq SU(4)_R$.

Recently there has been extensive analysis of such four point correlation functions, primarily for the simplest $p = 2$ case, in relation to the operator product expansion [1,2,3,4,5,6,7]. The operators appearing belong to various supermultiplets. For a long supermultiplet, for which the number of states is proportional to 2^{16}, the scale dimension Δ for the lowest dimension state is unconstrained so long as it satisfies an inequality $\Delta \geq \Delta_0$, where Δ_0 depends on the spin and $SU(4)_R$ symmetry representation, necessary for unitarity [8,9], for a useful review see [10]. In such supermultiplets the scale dimension ranges from Δ to $\Delta + 8$. The cardinal example [11] is the supermultiplet for which the lowest dimension state is formed by the Konishi scalar, which is a $SU(4)_R$ singlet and for which $\Delta_0 = 2$. In the interacting theory there is an anomalous dimension so that $\Delta = \Delta_0 + \mathrm{O}(\lambda)$. In addition there are various short supermultiplets in which the number of states involve factors 2^p, $p < 16$. These result from the lowest dimension state satisfying BPS like conditions where various supercharges annihilate the lowest dimension state. The simplest are the supermultiplets formed from the chiral primary operators described above which are scalars belonging to the $[0, p, 0]$ $SU(4)_R$ representation and with $\Delta_{\min} = p$. These are $\frac{1}{2}$-BPS representations and the supermultiplet then has operators with a maximum dimension $\Delta_{\min} + 4$, for $p \geq 4$. As well as this example the known short supermultiplets also include $\frac{1}{4}$-BPS representations with the lowest dimension states belonging to the $[q, p, q]$ $SU(4)_R$ representation $\Delta_{\min} = 2q + p$ and $\Delta_{\max} = \Delta_{\min} + 6$. For these examples the BPS shortening conditions apply to both the Q and $\bar{Q}$ supercharges which requires the lowest dimension state to be spinless. The scale dimensions of such short multiplets are expected to be protected against perturbative corrections.

Besides the contributions of these operators and their higher dimension descendents other operators with apparently protected scale dimensions have been identified in the operator product expansion analysis. The first to be exhibited [2,3] was a scalar operator

with $\Delta = 4$ in the 20-dimensional $[0,2,0]$ representation. This exactly satisfies the unitarity bound so there is no apparent reason to prohibit an anomalous dimension. This operator has been identified with the double trace operator formed from the product of two $[0,2,0]$ chiral primary operators in the large N limit and its vanishing anomalous dimension confirmed by field theory calculations [11,12,13]. Multiplets with protected dimensions, and with the possibility of non zero spins, were also demonstrated in the general tensor product decomposition of two chiral primary $\frac{1}{2}$-BPS operators by an analysis of the corresponding three point functions for $\mathcal{N} = 2$ [14] and $\mathcal{N} = 4$ [6]. Recently [7] we also showed the necessary existence in the operator expansion for two $[0,2,0]$ chiral primary operators of families of such protected supermultiplets whose lowest dimension operators satisfied $\Delta_0 = 4 + \ell$ where the spin representation was $(\frac{1}{2}\ell, \frac{1}{2}\ell)$ and which belonged to the $[0,2,0]$, $[1,0,1]$ representations for ℓ even, odd. The conformal partial wave analysis in [7] demonstrates that the associated supermultiplets do not have the full range of scale dimensions, and also spins, expected for a long supermultiplet.

In this paper we reanalyse the possible shortening conditions consistent with superconformal symmetry directly from the superconformal algebra and attempt to provide a unified treatment of different possible cases. We reproduce well known results for finding short supermultiplets when the lowest dimension operator satisfies BPS like conditions. If the BPS conditions are applied for both the Q and $\bar{Q}$ supercharges then the lowest dimension operator must be spinless, $j = \bar{j} = 0$. There are also various so called semi-short supermultiplets, depending on the $SU(4)_R$ representation of the lowest dimension state, which may occur at the threshold of the unitarity bound, $\Delta = \Delta_0$. In these examples the semi-shortening conditions may be applied for both the Q and $\bar{Q}$ supercharges for arbitrary spin representations $(j, \bar{j})$ and then the maximum scale dimension in the supermultiplets is $\Delta_0 + n$, $n = 4, 5, 6, 7$ with the total number of states involving a factor 2^{2n}. We also consider multiplets in which the semi-shortening condition is applied to the Q supercharges while for $\bar{Q}$ there is the full BPS shortening condition, and vice versa. The lowest dimension state then has spin $(j, 0)$ or $(0, \bar{j})$.

The structure of superconformal multiplets and possible shortening conditions has been extensively investigated previously by a variety of methods, in particular by using harmonic oscillator methods [15,16] and harmonic superspace [17,18,19]. Here we construct the supermultiplets directly from the superconformal algebra and also determine the spins and R-symmetry representations for the various fields present in supermultiplet which may contribute in a conformal operator product expansion. We attempt to give a unified description of possible short and semi-short multiplets including special cases where there are conservation equations or equations of motion. The particular cases depend on the fraction of the total number of supercharges for which the shortening or semi-shortening

conditions are applied. We also show how a long multiplet at its unitarity threshold may be decomposed in general into four semi-short multiplets although for a spinless lowest dimension state one is a $\frac{1}{4}$-BPS multiplet.

In $\mathcal{N} = 4$ supersymmetric gauge theories, except for $\frac{1}{2}$-BPS multiplets and one particular case for $\frac{1}{4}$-BPS multiplets, then even if fields are part of short or semi-short multiplets in the free theory there is no guarantee that there could not be a non zero anomalous dimension in the interacting theory for non zero g since they may all be components of a long supermultiplet. It is a non trivial exercise, not attempted here, to see how the spectrum of free field operators may be combined into different supermultiplets which may then be used to construct possible long multiplets. For the case of the Konishi scalar then the associated $\mathcal{N} = 1$ superfield satisfies a differential constraint in the free theory, corresponding to a semi-shortening condition on the supermultiplet, which is related to the presence of a conserved current. However this superspace equation has an anomaly in the interacting theory, which implies the usual axial anomaly for the conserved current, and the addition of the anomaly ensures that there are then sufficient degrees of freedom to form a long multiplet which may gain an anomalous dimension.

In our discussion an essential role is played by the Racah Speiser algorithm [20] for determining the different representations that may be formed when the supercharges act on a superconformal primary state. In general the corresponding Dynkin labels are obtained by adding the weights for the various supercharges to the Dynkin label for the superconformal primary state, although if one or more of the resulting Dynkin indices obtained in this way is negative then either the representation is replaced, up to a sign, by one with a Dynkin label with all indices positive or zero as usual or, as when one of the indices is -1, the associated representation is set to zero and is absent. This prescription allows a straightforward unification of the results for the different semi-shortening conditions. It also allows an easy understanding of when the fields satisfy conservation equations or equations of motion. A simple introduction to the Racah Speiser algorithm is given in an appendix. In special cases with this prescription a semi-short multiplet reduces to a short multiplet.

In detail in this paper in the next section we review the conformal algebra and describe briefly how unitary positive energy representations are constructed from conformal primary states. In particular we relate the states formed by the action of field operators on the vacuum to the finite norm positive energy states, as defined by an operator H, in the usual mathematical treatment. The eigenvalues Δ of H correspond to the scale dimensions of the fields as given by the dilation operator D (although D is hermitian it has eigenvalues $i\Delta$). In section 3 we review the superconformal algebra in four dimensions and relate the generators of the $U(2)_R$ and $SU(4)_R$ R-symmetry groups for $\mathcal{N} = 2$ and $\mathcal{N} = 4$

to those in the standard basis associated with simple roots. In section 4 the $\mathcal{N} = 2$ supermultiplets and the corresponding BPS and semi-shortening conditions analysed. It is shown how constraints such as conservation equations arise from the supersymmetry algebra. The $SU(2)_R$ and spin representations which arise in various supermultiplets are described pictorially. The case when a long multiplet can be decomposed as a semi-direct sum into short multiplets is described. The same analysis is repeated in section 5 for the $\mathcal{N} = 4$ case although there is a greater range of possibilities due to there being BPS conditions for a fraction $s = \frac{1}{4}, \frac{1}{2}, \frac{3}{4}, 1$ of the supercharges. There is a similar set of cases for the semi-shortening conditions and these can be applied independently to the action of the Q and $\bar{Q}$ supercharges. A general long multiplet at the unitarity threshold is decomposed into semi-short multiplets, if the initial state is spinless this includes a $\frac{1}{4}$-BPS multiplet. In section 6 we discuss some special cases which arise for small representations when the Racah Speiser algorithm gives rise to negative contributions in the multiplet. The presence of such terms is identified with the requirement for imposing equations of motion or conservation equations on particular fields. We relate the results obtained here to specific supermultiplets which have been found by different methods previously. Finally in section 7 we discuss the implications of our results for the spectrum of operators in $\mathcal{N} = 4$ supersymmetric gauge theories, in particular when operators which belong to a short or semi-short multiplet may be expected to have protected scale dimensions. Some details are deferred to three appendices. In appendix A we show how the usual two point function may be obtained algebraically using essentially just group theory. In appendix B we describe the Racah Speiser algorithm for decomposing tensor products while appendix C contains some tables of representations for short and semi-short multiplets.

2. Conformal Algebra

In d dimensions the conformal algebra is well known, for $\eta_{ab} = \text{diag.}(-1, 1 \ldots, 1)$ where $a, b = 0, 1, \ldots d-1$,

$$\begin{aligned}
[M_{ab}, P_c] &= i(\eta_{ac} P_b - \eta_{bc} P_a)\,, \qquad [M_{ab}, K_c] = i(\eta_{ac} K_b - \eta_{bc} K_a)\,, \\
[M_{ab}, M_{cd}] &= i(\eta_{ac} M_{bd} - \eta_{bc} M_{ad} - \eta_{ad} M_{bc} + \eta_{bd} M_{ac})\,, \\
[D, P_a] &= i P_a\,, \quad [D, K_a] = -i K_a\,, \qquad [K_a, P_b] = -2i M_{ab} - 2i\eta_{ab} D\,,
\end{aligned} \tag{2.1}$$

where P_a, K_a are the generators of translations, special conformal transformations, $M_{ab} = -M_{ba}$ are the generators for $SO(d-1, 1)$ and D is the generator of scale transformations and other commutators not shown are zero. The algebra corresponds to that for $SO(d, 2)$

since defining M_{AB}, $A, B = 0, 1, \ldots, d+1$ by

$$M_{AB} = \begin{pmatrix} M_{ab} & -\frac{1}{2}(P_a - K_a) & -\frac{1}{2}(P_a + K_a) \\ \frac{1}{2}(P_b - K_b) & 0 & D \\ \frac{1}{2}(P_b + K_b) & -D & 0 \end{pmatrix}, \tag{2.2}$$

then (2.1) is summarised by

$$[M_{AB}, M_{CD}] = i(\eta_{AC} M_{BD} - \eta_{BC} M_{AD} - \eta_{AD} M_{BC} + \eta_{BD} M_{AC}), \tag{2.3}$$

with $\eta_{AB} = \text{diag.}(-1, 1 \ldots, 1, -1)$. For physical applications we require unitary, so that ${M_{AB}}^\dagger = M_{AB}$, positive energy, where the energy is determined by

$$H = M_{0\,d+1} = -\tfrac{1}{2}(P_0 + K_0), \tag{2.4}$$

representations.

Acting on a multi-component quasi-primary field $\mathcal{O}_I(x)$ we have

$$\begin{aligned} [P_a, \mathcal{O}_I(x)] &= i\partial_a \mathcal{O}_I(x), \qquad [D, \mathcal{O}_I(x)] = i(x{\cdot}\partial + \Delta)\mathcal{O}_I(x), \\ [M_{ab}, \mathcal{O}_I(x)] &= i(x_a \partial_b - x_b \partial_a)\mathcal{O}_I(x) + \mathcal{O}_J(x){(s_{ab})^J}_I, \\ [K_a, \mathcal{O}_I(x)] &= i(x^2 \partial_a - 2x_a\, x{\cdot}\partial - 2\Delta x_a)\mathcal{O}_I(x) - 2\mathcal{O}_J(x){(s_{ab})^J}_I x^b, \end{aligned} \tag{2.5}$$

where ${(s_{ab})^I}_J$ are the appropriate finite dimensional spin matrices, obeying the algebra of M_{ab}, and Δ is the scale dimension. For such a quasi-primary field we define

$$|\mathcal{O}\rangle_I = \mathcal{O}_I(0)|0\rangle \tag{2.6}$$

as a set of conformal primary states satisfying

$$K_a|\mathcal{O}\rangle_I = 0, \qquad D|\mathcal{O}\rangle_I = i\Delta|\mathcal{O}\rangle_I, \qquad M_{ab}|\mathcal{O}\rangle_I = |\mathcal{O}\rangle_J {(s_{ab})^J}_I. \tag{2.7}$$

The space of states, on which the conformal representation is defined, is then spanned by vectors of the form

$$\prod_n P_{a_n}|\mathcal{O}\rangle_I. \tag{2.8}$$

For the conjugate field given by $\bar{\mathcal{O}}_{\bar{I}}(x) = \mathcal{O}_I(x)^\dagger$ we may also define

$${}_{\bar{I}}\langle\bar{\mathcal{O}}| = \langle 0|\bar{\mathcal{O}}_{\bar{I}}(0). \tag{2.9}$$

which satisfies

$${}_{\bar{I}}\langle\bar{\mathcal{O}}|K_a = 0, \qquad {}_{\bar{I}}\langle\bar{\mathcal{O}}|D = -i\Delta\, {}_{\bar{I}}\langle\bar{\mathcal{O}}|, \qquad {}_{\bar{I}}\langle\bar{\mathcal{O}}|M_{ab} = {(\bar{s}_{ab})_{\bar{I}}}^{\bar{J}}\, {}_{\bar{J}}\langle\bar{\mathcal{O}}|, \quad {(\bar{s}_{ab})_{\bar{I}}}^{\bar{J}} = {(s_{ab})^J}_I{}^*. \tag{2.10}$$

The relation to the standard treatment of unitary positive energy representations is obtained by a similarity transformation. With $\hat{a} = 1, \ldots d-1$,

$$\begin{aligned} -e^{-\frac{\pi}{2}M_{0d}} iD e^{\frac{\pi}{2}M_{0d}} &= H\,, \qquad e^{-\frac{\pi}{2}M_{0d}}(M_{\hat{a}\hat{b}}, -iM_{0\hat{a}})e^{\frac{\pi}{2}M_{0d}} = (M_{\hat{a}\hat{b}}, M_{d\hat{a}})\,, \\ e^{-\frac{\pi}{2}M_{0d}}(P_{\hat{a}}, -iP_0)e^{\frac{\pi}{2}M_{0d}} &= (\mathcal{E}^+_{\ \hat{a}}, \mathcal{E}^+_{\ d})\,, \qquad e^{-\frac{\pi}{2}M_{0d}}(K_{\hat{a}}, -iK_0)e^{\frac{\pi}{2}M_{0d}} = (\mathcal{E}^-_{\ \hat{a}}, \mathcal{E}^-_{\ d})\,. \end{aligned} \tag{2.11}$$

for H given by (2.4) and where

$$\mathcal{E}^\pm_{\ r} = M_{d+1\,r} \pm iM_{0\,r}\,, \quad r = 1, \ldots d\,, \qquad \mathcal{E}^-_{\ r} = \mathcal{E}^{+\,\dagger}_{\ r}\,. \tag{2.12}$$

These have the commutators

$$[\mathcal{E}^-_{\ r}, \mathcal{E}^+_{\ s}] = 2\delta_{rs}H - 2iM_{rs}\,, \quad [H, \mathcal{E}^\pm_{\ r}] = \pm\mathcal{E}^\pm_{\ s}\,, \quad [\mathcal{E}^+_{\ r}, \mathcal{E}^+_{\ s}] = 0\,, \tag{2.13}$$

with M_{rs} generators for the compact $SO(d)$ subgroup given by (2.3) with $\eta_{rs} = \delta_{rs}$.

If we now define

$$|\Delta\rangle_I = e^{-\frac{\pi}{2}M_{0d}}|\mathcal{O}\rangle_I\,, \qquad {}_{\bar{I}}\langle\Delta| = {}_{\bar{I}}\langle\bar{\mathcal{O}}|e^{-\frac{\pi}{2}M_{0d}} \tag{2.14}$$

then from (2.7),(2.10) and (2.11) it is easy to see that

$$\begin{aligned} &H|\Delta\rangle_I = \Delta|\Delta\rangle_I\,, \quad \mathcal{E}^-_{\ i}|\Delta\rangle_I = 0\,, \quad M_{rs}|\Delta\rangle_I = |\Delta\rangle_J(\hat{s}_{rs})^J{}_I\,, \\ &{}_{\bar{I}}\langle\Delta|H = \Delta\,{}_{\bar{I}}\langle\Delta|\,, \quad {}_{\bar{I}}\langle\Delta|\mathcal{E}^+_{\ i} = 0\,, \quad {}_{\bar{I}}\langle\Delta|M_{rs} = (\hat{\bar{s}}_{rs})_{\bar{I}}{}^{\bar{J}}\,{}_{\bar{J}}\langle\Delta|\,, \end{aligned} \tag{2.15}$$

where $\hat{s}_{rs}$ and $\hat{\bar{s}}_{rs}$ are matrices representing M_{rs} given by

$$\hat{s}_{\hat{a}\hat{b}} = s_{\hat{a}\hat{b}}\,, \quad \hat{s}_{d\hat{a}} = -is_{0\hat{a}}\,, \qquad \hat{\bar{s}}_{\hat{a}\hat{b}} = \bar{s}_{\hat{a}\hat{b}}\,, \quad \hat{\bar{s}}_{d\hat{a}} = i\bar{s}_{0\hat{a}}\,. \tag{2.16}$$

Corresponding to (2.8) we have a basis for a positive energy representation space given by $\prod_n \mathcal{E}^+_{\ r_n}|\Delta\rangle_I$. These states are normalisable,

$${}_{\bar{I}}\langle\Delta|\Delta\rangle_I = N_{\bar{I}I}\,, \tag{2.17}$$

where $N = [N_{\bar{I}I}]$ is a positive definite matrix satisfying

$$N^{-1}\hat{\bar{s}}_{rs}N = \hat{s}_{rs}\,. \tag{2.18}$$

The corresponding states defined in (2.6) and (2.9) are not normaliseable, unless $\mathcal{O}$ is the identity and $\Delta = 0$. The lack of a finite norm for $|\mathcal{O}\rangle_I$ is of course necessary for the hermitian operator D to have imaginary eigenvalues in (2.7). For subsequent discussion it is more convenient to use the non-normalisable states formed by the action of quantum fields at $x = 0$ although this is entirely equivalent to a discussion in terms of states with finite norm. In appendix A we show how the standard two point function for the field operators can be recovered algebraically from (2.17) using (2.15).

3. Superconformal Algebra, Four Dimensions

As is well known the conformal group in four dimensions may be extended by including supercharges $Q^i{}_\alpha$, $\bar{Q}_{i\dot{\alpha}}$ and also $S_i{}^\alpha$, $\bar{S}^{i\dot{\alpha}}$, $i = 1, \ldots \mathcal{N}$, so as to realise the Lie superalgebra for $SU(2,2|\mathcal{N})$. The usual supercharges satisfy

$$\{Q^i{}_\alpha, \bar{Q}_{j\dot{\alpha}}\} = 2\delta^i{}_j \mathrm{P}_{\alpha\dot{\alpha}}\,, \qquad \{Q^i{}_\alpha, Q^j{}_\beta\} = \{\bar{Q}_{i\dot{\alpha}}, \bar{Q}_{j\dot{\beta}}\} = 0\,, \tag{3.1}$$

and their superconformal extensions

$$\{\bar{S}^{i\dot{\alpha}}, S_j{}^\alpha\} = 2\delta^i{}_j \tilde{\mathrm{K}}^{\dot{\alpha}\alpha}\,, \qquad \{\bar{S}^{i\dot{\alpha}}, \bar{S}^{j\dot{\beta}}\} = \{S_i^\alpha, S_j^\beta\} = 0\,, \tag{3.2}$$

while the anti-commutators of the Q's and the S's are

$$\{Q^i{}_\alpha, \bar{S}^{j\dot{\alpha}}\} = 0\,, \qquad \{S_i^\alpha, \bar{Q}_{j\dot{\alpha}}\} = 0\,, \tag{3.3}$$

and also

$$\{Q^i{}_\alpha, S_j^\beta\} = 4\big(\delta^i{}_j (M_\alpha{}^\beta - \tfrac{1}{2} i\, \delta_\alpha{}^\beta D) - \delta_\alpha{}^\beta R^i{}_j\big)\,, \tag{3.4a}$$

$$\{\bar{S}^{i\dot{\alpha}}, \bar{Q}_{j\dot{\beta}}\} = 4\big(\delta^i{}_j (\bar{M}^{\dot{\alpha}}{}_{\dot{\beta}} + \tfrac{1}{2} i\, \delta^{\dot{\alpha}}{}_{\dot{\beta}} D) - \delta^{\dot{\alpha}}{}_{\dot{\beta}} R^i{}_j\big)\,, \tag{3.4b}$$

where

$$\begin{aligned} &\mathrm{P}_{\alpha\dot{\alpha}} = (\sigma^a)_{\alpha\dot{\alpha}} P_a\,, && \tilde{\mathrm{K}}^{\dot{\alpha}\alpha} = (\bar{\sigma}^a)^{\dot{\alpha}\alpha} K_a\,, \\ &M_\alpha{}^\beta = -\tfrac{1}{4} i (\sigma^a \bar{\sigma}^b)_\alpha{}^\beta M_{ab}\,, && \bar{M}^{\dot{\alpha}}{}_{\dot{\beta}} = -\tfrac{1}{4} i (\bar{\sigma}^a \sigma^b)^{\dot{\alpha}}{}_{\dot{\beta}} M_{ab}\,. \end{aligned} \tag{3.5}$$

In this basis the commutators for M_{ab} reduce to

$$[M_\alpha{}^\beta, M_\gamma{}^\delta] = \delta_\gamma{}^\beta M_\alpha{}^\delta - \delta_\alpha{}^\delta M_\gamma{}^\beta\,, \qquad [\bar{M}^{\dot{\alpha}}{}_{\dot{\beta}}, \bar{M}^{\dot{\gamma}}{}_{\dot{\delta}}] = -\delta^{\dot{\alpha}}{}_{\dot{\delta}} \bar{M}^{\dot{\gamma}}{}_{\dot{\beta}} + \delta^{\dot{\gamma}}{}_{\dot{\beta}} \bar{M}^{\dot{\alpha}}{}_{\dot{\delta}}\,, \tag{3.6}$$

and acting on the supercharges we have

$$\begin{aligned} &[M_\alpha{}^\beta, Q^i{}_\gamma] = \delta_\gamma{}^\beta Q^i{}_\alpha - \tfrac{1}{2}\delta_\alpha{}^\beta Q^i{}_\gamma\,, && [M_\alpha{}^\beta, S_i^\gamma] = -\delta_\alpha{}^\gamma S_i^\beta + \tfrac{1}{2}\delta_\alpha{}^\beta S_i^\gamma\,, \\ &[\bar{M}^{\dot{\alpha}}{}_{\dot{\beta}}, \bar{Q}_{i\dot{\gamma}}] = -\delta^{\dot{\alpha}}{}_{\dot{\gamma}} \bar{Q}_{i\dot{\beta}} + \tfrac{1}{2}\delta^{\dot{\alpha}}{}_{\dot{\beta}} \bar{Q}_{i\dot{\gamma}}\,, && [\bar{M}^{\dot{\alpha}}{}_{\dot{\beta}}, \bar{S}^{i\dot{\gamma}}] = \delta^{\dot{\gamma}}{}_{\dot{\beta}} \bar{S}^{i\dot{\alpha}} - \tfrac{1}{2}\delta^{\dot{\alpha}}{}_{\dot{\beta}} \bar{S}^{i\dot{\gamma}}\,. \end{aligned} \tag{3.7}$$

Under the action of the generator of scale transformations we have

$$[D, Q^i{}_\alpha] = \tfrac{1}{2} i Q^i{}_\alpha\,, \quad [D, \bar{Q}_{i\dot{\alpha}}] = \tfrac{1}{2} i \bar{Q}_{i\dot{\alpha}}\,, \qquad [D, S_i^\alpha] = -\tfrac{1}{2} i S_i^\alpha\,, \quad [D, \bar{S}^{i\dot{\alpha}}] = -\tfrac{1}{2} i \bar{S}^{i\dot{\alpha}}\,, \tag{3.8}$$

and also we have

$$[K_a, Q^i{}_\alpha] = -(\sigma_a)_{\alpha\dot{\alpha}} \bar{S}^{i\dot{\alpha}}\,, \qquad [K_a, \bar{Q}_{i\dot{\alpha}}] = S_i^\alpha (\sigma_a)_{\alpha\dot{\alpha}}\,, \tag{3.9a}$$

$$[P_a, \bar{S}^{i\dot{\alpha}}] = -(\bar{\sigma}_a)^{\dot{\alpha}\alpha} Q^i{}_\alpha\,, \qquad [P_a, S_i^\alpha] = \bar{Q}_{i\dot{\alpha}} (\bar{\sigma}_a)^{\dot{\alpha}\alpha}\,. \tag{3.9b}$$

The remaining generators involve the $U(\mathcal{N})$ R-symmetry for which the Lie algebra is

$$[R^i{}_j, R^k{}_l] = \delta^k{}_j R^i{}_l - \delta^i{}_l R^k{}_j \,. \tag{3.10}$$

These act on the supercharges according to

$$\begin{aligned} [R^i{}_j, Q^k{}_\alpha] &= \delta^k{}_j Q^i{}_\alpha - \tfrac{1}{4}\delta^i{}_j Q^k{}_\alpha \,, \qquad & [R^i{}_j, \bar{Q}_{k\dot\alpha}] &= -\delta^i{}_k \bar{Q}_{j\dot\alpha} + \tfrac{1}{4}\delta^i{}_j \bar{Q}_{k\dot\alpha} \,, \\ [R^i{}_j, S_k{}^\alpha] &= -\delta^i{}_k S_j{}^\alpha + \tfrac{1}{4}\delta^i{}_j S_k{}^\alpha \,, \qquad & [R^i{}_j, \bar{S}^{k\dot\alpha}] &= \delta^k{}_j \bar{S}^{i\dot\alpha} - \tfrac{1}{4}\delta^i{}_j \bar{S}^{k\dot\alpha} \,. \end{aligned} \tag{3.11}$$

For $\mathcal{N} = 4$ it is evident that we may impose $R^i_i = 0$ so that the R-symmetry is then $SU(4)$ and the supergroup reduces to $PSU(2,2|4)$.

As operators we also require the hermeticity conditions

$$Q^i{}_\alpha{}^\dagger = \bar{Q}_{i\dot\alpha} \,, \quad S_i^\alpha{}^\dagger = \bar{S}^{i\dot\alpha} \,, \quad M_\alpha{}^{\beta\dagger} = \bar{M}^{\dot\beta}{}_{\dot\alpha} \,, \quad R^i{}_j{}^\dagger = R^j{}_i \,. \tag{3.12}$$

Applying the transformation (2.11) to the supercharges, using $(3.9a,b)$, we define

$$\begin{aligned} \mathcal{Q}^{+i}{}_\alpha &= e^{-\frac{\pi}{2}M_{0d}} Q^i{}_\alpha e^{\frac{\pi}{2}M_{0d}} = \tfrac{1}{\sqrt{2}}\big(Q^i{}_\alpha + \sigma_{0\,\alpha\dot\alpha}\bar{S}^{i\dot\alpha}\big) \,, \\ \mathcal{Q}^{-\alpha}_i &= e^{-\frac{\pi}{2}M_{0d}} S_i^\alpha e^{\frac{\pi}{2}M_{0d}} = \tfrac{1}{\sqrt{2}}\big(S_i^\alpha + \bar{Q}_{i\dot\alpha}\,\bar\sigma_0{}^{\dot\alpha\alpha}\big) \,, \\ -\mathcal{S}^{-i\dot\alpha} &= e^{-\frac{\pi}{2}M_{0d}} \bar{S}^{i\dot\alpha} e^{\frac{\pi}{2}M_{0d}} = \tfrac{1}{\sqrt{2}}\big(\bar{S}^{i\dot\alpha} - \bar\sigma_0{}^{\dot\alpha\alpha} Q^i{}_\alpha\big) \,, \\ \mathcal{S}^+_{i\dot\alpha} &= e^{-\frac{\pi}{2}M_{0d}} \bar{Q}_{i\dot\alpha} e^{\frac{\pi}{2}M_{0d}} = \tfrac{1}{\sqrt{2}}\big(\bar{Q}_{i\dot\alpha} - S_i^\alpha \sigma_{0\,\alpha\dot\alpha}\big) \,. \end{aligned} \tag{3.13}$$

and the algebra $(3.4a,b)$ becomes

$$\begin{aligned} \{\mathcal{Q}^{+i}{}_\alpha, \mathcal{Q}^{-\beta}_j\} &= 2\,\delta^i{}_j \delta_\alpha{}^\beta H + 4\,\delta^i{}_j \tilde{M}_\alpha{}^\beta - 4\,\delta_\alpha{}^\beta R^i{}_j \,, \\ \{\mathcal{S}^{-i\dot\alpha}, \mathcal{S}^+_{j\dot\beta}\} &= 2\,\delta^i{}_j \delta^{\dot\alpha}{}_{\dot\beta} H - 4\,\delta^i{}_j \tilde{\bar{M}}^{\dot\alpha}{}_{\dot\beta} + 4\,\delta^{\dot\alpha}{}_{\dot\beta} R^i{}_j \,, \end{aligned} \tag{3.14}$$

for $\tilde{M}_\alpha{}^\beta = e^{-\frac{\pi}{2}M_{0d}} M_\alpha{}^\beta e^{\frac{\pi}{2}M_{0d}}$, $\tilde{\bar{M}}^{\dot\alpha}{}_{\dot\beta} = e^{-\frac{\pi}{2}M_{0d}} \bar{M}^{\dot\alpha}{}_{\dot\beta} e^{\frac{\pi}{2}M_{0d}}$. In addition we have

$$\begin{aligned} \{\mathcal{Q}^{+i}{}_\alpha, \mathcal{S}^+_{j\dot\alpha}\} &= 2\,\delta^i{}_j (e_r)_{\alpha\dot\alpha} \mathcal{E}^+{}_r \,, \qquad & e_r &= (\sigma_{\hat a}, -i\sigma_0) \,, \\ \{\mathcal{S}^{-i\dot\alpha}, \mathcal{Q}^{-\alpha}_j\} &= 2\,\delta^i{}_j (\bar{e}_r)^{\dot\alpha\alpha} \mathcal{E}^-{}_r \,, \qquad & \bar{e}_r &= (-\bar\sigma_{\hat a}, i\bar\sigma_0) \,, \end{aligned} \tag{3.15}$$

and $[H, \mathcal{Q}^\pm] = \pm\frac{1}{2}\mathcal{Q}^\pm$, $[H, \mathcal{S}^\pm]] = \pm\frac{1}{2}\mathcal{S}^\pm$. From the hermeticity condition (3.12)

$$(\mathcal{Q}^{+i}{}_\alpha)^\dagger = \mathcal{Q}^{-\beta}_i \sigma_{0\beta\dot\alpha} \,, \qquad (\mathcal{S}^+_{i\dot\alpha})^\dagger = \sigma_{0\alpha\dot\beta} \mathcal{S}^{-i\dot\beta} \,, \tag{3.16}$$

and also $(\tilde{M}_\alpha{}^\beta)^\dagger = \bar\sigma_0{}^{\dot\beta\gamma} \tilde{M}_\gamma{}^\delta \sigma_{0\delta\dot\alpha}$. With the conventional choice $\sigma_0 = \bar\sigma_0 = 1$ it is clear that the chiral subalgebra given by $\{\mathcal{Q}^{+i}{}_\alpha, \mathcal{Q}^{-\alpha}_i, \tilde{M}_\alpha{}^\beta, R^i{}_j, H\}$, or $\{\mathcal{S}^{-i\dot\alpha}, \mathcal{S}^+_{i\dot\alpha}, \tilde{\bar{M}}^{\dot\alpha}{}_{\dot\beta}, R^i{}_j, H\}$, are generators for $SU(2|4)$ and H has a positive spectrum. Finding the unitary finite

dimensional representations of these subalgebras, with lowest energy states $|\Delta\rangle$ satisfying as well as (2.15) $\mathcal{Q}_i^{-\alpha}|\Delta\rangle = \mathcal{S}^{-i\dot{\alpha}}|\Delta\rangle = 0$, is a necessary preliminary subsequently for determining the unitary positive energy representations of the full superconformal group.

In this paper we are interested in discussing just the physical cases of $\mathcal{N} = 2$ and $\mathcal{N} = 4$. For $\mathcal{N} = 2$ we may simply write

$$[R^i{}_j] = \begin{pmatrix} R_3 & R_+ \\ R_- & -R_3 \end{pmatrix} + \tfrac{1}{2}\hat{R}\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \tag{3.17}$$

where $R_\pm, R_3$ form a standard $SU(2)$ algebra and $\hat{R}$ corresponds to the generator of $U(1)_R$. From (3.11) it is easy to see that $Q^i{}_\alpha$, $\bar{Q}_{i\dot{\alpha}}$ belong to $R = \frac{1}{2}$ representations since we have

$$\begin{aligned} &[R_+, Q^1{}_\alpha] = 0\,, \qquad [R_3, Q^1{}_\alpha] = \tfrac{1}{2}Q^1{}_\alpha\,, \qquad [R_-, Q^1{}_\alpha] = Q^2{}_\alpha\,, \\ &[R_+, \bar{Q}_{2\dot{\alpha}}] = 0\,, \qquad [R_3, \bar{Q}_{2\dot{\alpha}}] = \tfrac{1}{2}\bar{Q}_{2\dot{\alpha}}\,, \qquad [R_-, \bar{Q}_{2\dot{\alpha}}] = -\bar{Q}_{1\dot{\alpha}}\,. \end{aligned} \tag{3.18}$$

For $\mathcal{N} = 4$ we express $R^i{}_j$ in terms of the generators of $SU(4)$ in a Chevalley basis. For each simple root there is an associated $SU(2)$ algebra given by $E_i^\pm$, H_i, $i = 1, \ldots r$ so that r is the rank, which have the commutation relations

$$[H_i, H_j] = 0\,, \quad [E_i{}^+, E_j{}^-] = \delta_{ij}H_j\,, \quad [H_i, E_j{}^\pm] = \pm K_{ji}E_j{}^\pm\,, \text{ no sum on } j\,, \tag{3.19}$$

where K_{ji} are the elements of the Cartan matrix, $K_{ii} = 2$. The generators for the non simple positive roots are obtained by appropriate commutators of the E_i^+, and for the corresponding negative roots from the equivalent commutators of the E_i^-, subject to the Serre relations,

$$[\underbrace{E_i^\pm, [\ldots [E_i{}^\pm}_{1-K_{ji}}, E_j{}^\pm]\ldots]] = 0\,, \qquad i \neq j\,. \tag{3.20}$$

The remaining commutators for any pair of generators are then determined using (3.19) and the Jacobi identity. For a compact group we may further impose the hermeticity conditions

$$H_i{}^\dagger = H_i\,, \qquad E_i{}^{+\,\dagger} = E_i^-\,, \qquad i = 1, \ldots, r\,. \tag{3.21}$$

For any representation space a convenient basis is then given by the eigenvectors of H_i

$$H_i|\lambda_1, \lambda_2, \ldots\rangle = \lambda_i|\lambda_1, \lambda_2, \ldots\rangle\,, \tag{3.22}$$

where the λ_i take integer values. The representation is then uniquely characterised by the highest weight vector satisfying

$$E_i^+|\lambda_1, \lambda_2, \ldots\rangle^{\rm hw} = 0\,, \qquad \lambda_i \geq 0\,, \tag{3.23}$$

and the representation then has Dynkin labels $[\lambda_1, \lambda_2, \ldots]$. The remaining basis vectors $|\lambda_1, \lambda_2, \ldots\rangle$ for the representation are then obtained by the successive action of E_i^- on the highest weight vector. We may also note that

$$E_i^-|\lambda_1, \lambda_2, \ldots\rangle^{\rm hw} = 0 \qquad \text{if} \qquad \lambda_i = 0\,. \tag{3.24}$$

For $SU(4)$, of rank 3, the Cartan matrix is

$$[K_{ij}] = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{pmatrix}\,. \tag{3.25}$$

For this case we may satisfy (3.10) by taking

$$[R^i{}_j] = \begin{pmatrix} \frac{1}{4}(3H_1{+}2H_2{+}H_3) & E_1{}^+ & [E_1{}^+, E_2{}^+] & [E_1{}^+, [E_2{}^+, E_3{}^+]] \\ E_1{}^- & \frac{1}{4}(-H_1{+}2H_2{+}H_3) & E_2{}^+ & [E_2{}^+, E_3{}^+] \\ -[E_1{}^-, E_2{}^-] & E_2{}^- & -\frac{1}{4}(H_1{+}2H_2{-}H_3) & E_3{}^+ \\ [E_1{}^-, [E_2{}^-, E_3{}^-]] & -[E_2{}^-, E_3{}^-] & E_3{}^- & -\frac{1}{4}(H_1{+}2H_2{+}3H_3) \end{pmatrix}\,. \tag{3.26}$$

All commutators follow from (3.19), (3.20) and the Jacobi identity. The supercharges $Q^i{}_\alpha$, $\bar{Q}_{i\dot\alpha}$ correspond to the $[1,0,0]$, $[0,0,1]$ representations and from (3.11) we then have

$$\begin{aligned} &[H_1, Q^1{}_\alpha] = Q^1{}_\alpha\,, \qquad [H_i, Q^1{}_\alpha] = 0\,,\ i = 2,3\,, \qquad [E_i{}^+, Q^1{}_\alpha] = 0\,, \\ &[H_3, \bar{Q}_{4\dot\alpha}] = \bar{Q}_{4\dot\alpha}\,, \qquad [H_i, \bar{Q}_{4\dot\alpha}] = 0\,,\ i = 1,2\,, \qquad [E_i{}^+, \bar{Q}_{4\dot\alpha}] = 0\,. \end{aligned} \tag{3.27}$$

Under commutation with E_i^- we have

$$Q^1{}_\alpha \xrightarrow{E_1{}^-} Q^2{}_\alpha \xrightarrow{E_2{}^-} Q^3{}_\alpha \xrightarrow{E_3{}^-} Q^4{}_\alpha\,, \qquad \bar{Q}_{4\dot\alpha} \xrightarrow{E_3{}^-} -\bar{Q}_{3\dot\alpha} \xrightarrow{E_2{}^-} \bar{Q}_{2\dot\alpha} \xrightarrow{E_1{}^-} -\bar{Q}_{1\dot\alpha}\,. \tag{3.28}$$

In general $SU(4)$ representations with Dynkin labels $[\lambda_1, \lambda_2, \lambda_3]$ have dimensions

$$d(\lambda_1, \lambda_2, \lambda_3) = \tfrac{1}{12}(\lambda_1{+}\lambda_2{+}\lambda_3{+}3)(\lambda_1{+}\lambda_2{+}2)(\lambda_2{+}\lambda_3{+}2)(\lambda_1{+}1)(\lambda_2{+}1)(\lambda_3{+}1)\,. \tag{3.29}$$

4. Superconformal Representations, $\mathcal{N} = 2$

We consider superconformal representations in which the lowest dimension states belong to a $SU(2)_R$ representation, with $U(1)_R$ charge r, which is formed by the action of R_- on a highest weight $SU(2)$ state $|R, r\rangle^{\rm hw}$ satisfying

$$R_+|R, r\rangle^{\rm hw} = 0\,, \qquad R_3|R, r\rangle^{\rm hw} = R|R, r\rangle^{\rm hw}\,, \qquad \hat{R}|R, r\rangle^{\rm hw} = r|R, r\rangle^{\rm hw}\,. \tag{4.1}$$

For the consideration of spin we here also adopt a spinorial basis for the $(j,\bar{j})$ representation given by

$$|R,r\rangle_{\alpha_1\ldots\alpha_{2j},\dot{\alpha}_1\ldots\dot{\alpha}_{2\bar{j}}} = |R,r\rangle_{(\alpha_1\ldots\alpha_{2j}),(\dot{\alpha}_1\ldots\dot{\alpha}_{2\bar{j}})}\,. \tag{4.2}$$

Under the action of the spin generators $M_\alpha{}^\beta$, $\bar{M}^{\dot{\beta}}{}_{\dot{\alpha}}$, defined in (3.5), the states given by (4.2) transform according to

$$\begin{aligned} M_\alpha{}^\beta|R,r\rangle_{\alpha_1\ldots\alpha_{2j},\dot{\alpha}_1\ldots\dot{\alpha}_{2\bar{j}}} &= j\big(2\delta_{(\alpha_1}{}^\beta|R,r\rangle_{\alpha_2\ldots\alpha_{2j})\alpha,\dot{\alpha}_1\ldots\dot{\alpha}_{2\bar{j}}} - \delta_\alpha{}^\beta|R,r\rangle_{\alpha_1\ldots\alpha_{2j},\dot{\alpha}_1\ldots\dot{\alpha}_{2\bar{j}}}\big)\,,\\ \bar{M}^{\dot{\beta}}{}_{\dot{\alpha}}|R,r\rangle_{\alpha_1\ldots\alpha_{2j},\dot{\alpha}_1\ldots\dot{\alpha}_{2\bar{j}}} &= -\bar{j}\big(2|R,r\rangle_{\alpha_1\ldots\alpha_{2j},\dot{\alpha}(\dot{\alpha}_1\ldots\dot{\alpha}_{2\bar{j}-1}}\delta^{\dot{\beta}}{}_{\dot{\alpha}_{2\bar{j}})} - \delta^{\dot{\beta}}{}_{\dot{\alpha}}|R,r\rangle_{\alpha_1\ldots\alpha_{2j},\dot{\alpha}_1\ldots\dot{\alpha}_{2\bar{j}}}\big)\,. \end{aligned} \tag{4.3}$$

The representation formed from states satisfying (4.1) and (4.3) is labelled $R_{(j,\bar{j})}$.

A highest weight conformal primary state is then required to satisfy from (2.7)

$$K_a|R,r\rangle^{\rm hw}_{\alpha_1\ldots\alpha_{2j},\dot{\alpha}_1\ldots\dot{\alpha}_{2\bar{j}}} = 0 \qquad D|R,r\rangle^{\rm hw}_{\alpha_1\ldots\alpha_{2j},\dot{\alpha}_1\ldots\dot{\alpha}_{2\bar{j}}} = i\Delta|R,r\rangle^{\rm hw}_{\alpha_1\ldots\alpha_{2j},\dot{\alpha}_1\ldots\dot{\alpha}_{2\bar{j}}}\,. \tag{4.4}$$

In addition from (3.8) we must impose

$$S_i^\alpha|R,r\rangle^{\rm hw}_{\alpha_1\ldots\alpha_{2j},\dot{\alpha}_1\ldots\dot{\alpha}_{2\bar{j}}} = \bar{S}^{i\dot{\alpha}}|R,r\rangle^{\rm hw}_{\alpha_1\ldots\alpha_{2j},\dot{\alpha}_1\ldots\dot{\alpha}_{2\bar{j}}} = 0\,, \tag{4.5}$$

for this to the lowest dimension in the supermultiplet and to be a superconformal primary state.

A basis for the full representation space is then provided by acting on states satisfying (4.4) and (4.5), as well as (4.1), with R_-, P_a and the eight supercharges $Q^i{}_\alpha$, $\bar{Q}_{k\dot{\alpha}}$. The set of states $\mathcal{S}_{l,\bar{l}}$

$$\prod_{i,k,\alpha,\dot{\alpha}} \big(Q^i{}_\alpha\big)^{n_{i\alpha}}\big(\bar{Q}_{k\dot{\alpha}}\big)^{\bar{n}_{k\dot{\alpha}}}|R,r\rangle^{\rm hw}_{\alpha_1\ldots\alpha_{2j},\dot{\alpha}_1\ldots\dot{\alpha}_{2\bar{j}}}\,, \qquad n_{i\alpha},\bar{n}_{k\dot{\alpha}} = 0,1\,, \tag{4.6}$$

for $l = \sum_{i,\alpha} n_{i\alpha}$, $\bar{l} = \sum_{k,\dot{\alpha}} \bar{n}_{k\dot{\alpha}}$, $l,\bar{l} = 0,1,2,3,4$, and for some appropriate ordering of the Q and $\bar{Q}$ supercharges, form, together with those obtained by the action of R_-, a basis for a vector space $\mathcal{V}_{l,\bar{l}}$. For all states in $\mathcal{V}_{l,\bar{l}}$ the scale dimension is $\Delta + \frac{1}{2}(l+\bar{l})$ while the $U(1)_R$ charge is $r + \frac{1}{2}(l-\bar{l})$. Acting with K_a, using (3.9a) and (3.4a,b), (3.3) with (4.4), (4.5), $\mathcal{V}_{l,\bar{l}} \to \mathcal{V}_{l-1,\bar{l}-1}$ and $K_a\mathcal{V}_{l,0} = 0$, $K_a\mathcal{V}_{0,\bar{l}} = 0$. For general Δ any state (4.6) in $\mathcal{S}_{l,\bar{l}}$ is a conformal primary state, annihilated by K_a, if $l = 0$ or $\bar{l} = 0$ or becomes a conformal primary by the addition of a suitable combination of states in $\mathcal{V}_{l-1,\bar{l}-1}$ acted on by the momentum operator. The states belonging to $\mathcal{V}_{l,\bar{l}}$ may be decomposed into representations for $SU(2)_R$ and spin. According to the Racah Speiser algorithm the resulting representations $R'_{(j',\bar{j}')}$ are obtained by adding the weights associated with each supercharge, from (3.18),

$$Q^1{}_\alpha \sim \tfrac{1}{2}{}_{(\pm\frac{1}{2},0)}\,, \quad Q^2{}_\alpha \sim (-\tfrac{1}{2})_{(\pm\frac{1}{2},0)}\,, \quad \bar{Q}_{2\dot{\alpha}} \sim \tfrac{1}{2}{}_{(0,\pm\frac{1}{2})}\,, \quad \bar{Q}_{1\dot{\alpha}} \sim (-\tfrac{1}{2})_{(0,\pm\frac{1}{2})}\,, \tag{4.7}$$

to $R_{(j,\bar{j})}$, so long as all $R', j', \bar{j}' \geq 0$, for all possible $n_{i\alpha}, \bar{n}_{k\dot{\alpha}}$ consistent with fixed $l, \bar{l}$. Thus the $SU(2)_R$ representations are then given by every $R' = R + \frac{1}{2}\sum_\alpha (n_{1\alpha} - n_{2\alpha}) - \frac{1}{2}\sum_{\dot{\alpha}}(\bar{n}_{1\dot{\alpha}} - \bar{n}_{2\dot{\alpha}})$. Altogether there are therefore in general $\binom{4}{l}\binom{4}{\bar{l}}$ representations $R'_{(j',\bar{j}')}$. If one or more of the resulting $R', j', \bar{j}'$ are negative then, according the Racah Speiser algorithm which is discussed more fully later, there is a simple recipe for cancelling such representations. The four $Q^i{}_\alpha$ supercharges by themselves acting on a representation $R_{(j,\bar{j})}$ give

$$R_{(j,\bar{j})} \xrightarrow{Q} \begin{matrix}(R+\frac{1}{2})_{(j\pm\frac{1}{2},\bar{j})} \\ (R-\frac{1}{2})_{(j\pm\frac{1}{2},\bar{j})}\end{matrix} \xrightarrow{Q^2} \begin{matrix}(R\pm 1)_{(j,\bar{j})}, R_{(j,\bar{j})} \\ R_{(j\pm1,\bar{j})}, R_{(j,\bar{j})}\end{matrix} \xrightarrow{Q^3} \begin{matrix}(R+\frac{1}{2})_{(j\pm\frac{1}{2},\bar{j})} \\ (R-\frac{1}{2})_{(j\pm\frac{1}{2},\bar{j})}\end{matrix} \xrightarrow{Q^4} R_{(j,\bar{j})}\ , \quad (4.8)$$

corresponding to $l = 0, 1, 2, 3, 4$, $\bar{l} = 0$, so long as $R \geq 1$, $j \geq 1$. The action of the $\bar{Q}$ supercharges is given by the conjugate of (4.8).

A general long supermultiplet, denoted by $\mathcal{A}^{\Delta}_{R,r(j,\bar{j})}$, is obtained by the application of all 8 supercharges $Q^i{}_\alpha$, $\bar{Q}_{i\dot{\alpha}}$ to a highest weight state satisfying (4.1) and with $U(2)_R$ and spin quantum numbers $R, r, j, \bar{j}$. The representations belonging to $\mathcal{V}_{l,0}$ are given by (4.8) and those forming $\mathcal{V}_{l,\bar{l}}$, $\bar{l} > 0$ may then be obtained by applying the conjugate of (4.8) for all representations in $\mathcal{V}_{l,0}$. Since there are no constraints it is straightforward to see that

$$\dim \mathcal{A}^{\Delta}_{R,r(j,\bar{j})} = 256(2R+1)(2j+1)(2\bar{j}+1)\,. \quad (4.9)$$

For a unitary representation we must have in this case [8]

$$\Delta \geq 2 + 2j + 2R + r\,,\ 2 + 2\bar{j} + 2R - r\,. \quad (4.10)$$

For a truncated representation we may impose the BPS-condition if $j = 0$,

$$Q^1{}_\alpha |R, r\rangle^{\rm hw}_{\dot{\alpha}_1 \ldots \dot{\alpha}_{2\bar{j}}} = 0\,. \quad (4.11)$$

In this case we must impose consistency with the result (3.4a) for the anti-commutator $\{Q^1{}_\alpha, S_j{}^\beta\}$ and with (4.1) and (4.4), since the state is annihilated by $M_\alpha{}^\beta$, it is easy to see that this requires

$$\Delta = 2R + r\,. \quad (4.12)$$

For the $\bar{Q}$ charges the corresponding condition, if $\bar{j} = 0$, is

$$\bar{Q}_{2\dot{\alpha}} |R, r\rangle^{\rm hw}_{\alpha_1 \ldots \alpha_{2j}} = 0\,, \quad (4.13)$$

which gives instead from (3.4b)

$$\Delta = 2R - r\,. \quad (4.14)$$

With (4.11) or (4.13) the multiplet is generated using only ${Q^2}_\alpha$ or $\bar{Q}_{1\dot{\alpha}}$ so that $l = 0, 1, 2$ or $\bar{l} = 0, 1, 2$. The following representations are then obtained for the action of the Q or $\bar{Q}$ supercharges,

$$R_{(0,\bar{\jmath})} \xrightarrow{Q} (R-\tfrac{1}{2})_{(\frac{1}{2},\bar{\jmath})} \xrightarrow{Q^2} (R-1)_{(0,\bar{\jmath})}\,, \qquad R_{(j,0)} \xrightarrow{\bar{Q}} (R-\tfrac{1}{2})_{(j,\frac{1}{2})} \xrightarrow{\bar{Q}^2} (R-1)_{(j,0)}\,, \tag{4.15}$$

with dimensions $8R(2\bar{\jmath}+1)$, $8R(2j+1)$ in each case.

If (4.11) holds for both ${Q^1}_\alpha$ and ${Q^2}_\alpha$ then it is necessary that $R = 0$ and $\Delta = r$. Conversely for $R = 0$ (4.11), using (3.18), implies the corresponding result for ${Q^2}_\alpha$.

Reduced supermultiplets may be obtained by applying shortening conditions to the action of the Q or $\bar{Q}$ supercharges or both. Imposing just (4.11), and hence requiring (4.12), we may construct an asymmetric supermultiplet, using the conjugate of (4.8), which is denoted by $\mathcal{B}_{R,r(0,\bar{\jmath})}$. This can then be represented by the diagram, corresponding to $l = 0, 1, 2,\ \bar{l} = 0, 1, 2, 3, 4$,



(4.16)

where $\swarrow$ corresponds to the action of the Q supercharges and $\searrow$ to $\bar{Q}$. The dimension in this case is given by

$$\dim \mathcal{B}_{R,r(0,\bar{\jmath})} = 128R(2\bar{\jmath}+1)\,. \tag{4.17}$$

Unitarity here requires

$$r \geq \bar{\jmath}+1\,. \tag{4.18}$$

We may similarly construct the conjugate supermultiplet $\bar{\mathcal{B}}_{R,r(j,0)}$ with, instead of (4.18), $-r \geq j+1$ and the lowest scale dimension given by (4.14) where

$$\dim \bar{\mathcal{B}}_{R,r(j,0)} = 128R(2j+1)\,. \tag{4.19}$$

For $R=0$, when the lowest state is annihilated by $Q^i{}_\alpha$ for both $i=1,2$ and the multiplet is generated solely by the action of the $\bar{Q}$ charges, we have a chiral multiplet $\mathcal{E}_{r(0,\bar{\jmath})}$



(4.20)

For unitarity we must have (4.18) in this case as well and

$$\dim \mathcal{E}_{r(0,\bar{\jmath})} = 16(2\bar{\jmath}+1)\,. \tag{4.21}$$

Of course there is a corresponding conjugate chiral multiplet $\bar{\mathcal{E}}_{r(j,0)}$.

Imposing the conditions (4.11) and (4.13) simultaneously requires $r=0$ and also $j=\bar{\jmath}=0$. We may describe this $\mathcal{N}=2$ short supermultiplet $\hat{\mathcal{B}}_R$, generated by the action

of Q^2, $\bar{Q}_1$ as in (4.15), by the diagram



(4.22)

It is easy to see that the total dimension of the representations in (4.22) is

$$\dim \hat{\mathcal{B}}_R = 16(2R-1)\,. \tag{4.23}$$

For $j > 0$ it is also possible to impose conditions which lead to multiplet shortening. We consider the condition corresponding to the absence of the representations with spin $(j-\frac{1}{2},\bar{\jmath})$ at the first level,

$$\varepsilon^{\alpha\beta} Q^i{}_\beta |R,r\rangle^{\rm hw}_{\alpha_1\ldots\alpha_{2j-1}\alpha,\dot\alpha_1\ldots\dot\alpha_{2\bar\jmath}} = 0\,, \tag{4.24}$$

The necessary consistency conditions then arise from $\varepsilon^{\alpha\beta}\{Q^i{}_\beta, S_l{}^\gamma\}|R,r\rangle^{\rm hw}_{\alpha_1\ldots\alpha_{2j-1}\alpha,\dot\alpha_1\ldots\dot\alpha_{2\bar\jmath}} = 0$ and we may easily find from (3.4a) and (4.3)

$$\big(\delta^i_l(-j-1+\tfrac{1}{2}\Delta) - R^i_l\big)|R,r\rangle^{\rm hw}_{\alpha_1\ldots\alpha_{2j},\dot\alpha_1\ldots\dot\alpha_{2\bar\jmath}} = 0\,, \qquad l=1,2\,. \tag{4.25}$$

If we require (4.24) just for $i=1$ then this gives

$$\Delta = 2 + 2j + 2R + r\,, \tag{4.26}$$

and the action of the Q supercharges on the highest weight state gives the representations

$$R_{(j,\bar\jmath)} \xrightarrow{Q} \begin{matrix}(R+\frac{1}{2})_{(j+\frac{1}{2},\bar\jmath)}\\ (R-\frac{1}{2})_{(j\pm\frac{1}{2},\bar\jmath)}\end{matrix} \xrightarrow{Q^2} \begin{matrix}R_{(j+1,\bar\jmath),(j,\bar\jmath)}\\ (R-1)_{(j,\bar\jmath)}\end{matrix} \xrightarrow{Q^3} (R-\tfrac{1}{2})_{(j+\frac{1}{2},\bar\jmath)}\,, \tag{4.27}$$

with a dimension $8(R(4j+3)+j+1)(2\bar\jmath+1)$. If (4.24) is imposed for both $i=1,2$ then (4.25) requires $R=0$ and $\Delta = 2+2j+r$. Conversely if $R=0$ the condition (4.24) for

$i=1$ implies also the $i=2$ case by application of R_-. Assuming (4.24) for $i=1,2$ the Q supercharges give for general R

$$R_{(j,\bar{\jmath})} \xrightarrow{Q} (R\pm\tfrac{1}{2})_{(j+\frac{1}{2},\bar{\jmath})} \xrightarrow{Q^2} R_{(j+1,\bar{\jmath})}\,. \tag{4.28}$$

The dimension of these representations is $8(2R+1)(j+1)(2\bar{\jmath}+1)$.

For $j=0$ the condition (4.24) is replaced by

$$(Q^i)^2|R,r\rangle^{\rm hw}_{\dot\alpha_1\ldots\dot\alpha_{2\bar\jmath}}=0\,. \tag{4.29}$$

The corresponding condition for $j>0$ is of course a consequence of (4.24). In this case conditions on representations appearing in the supermultiplet arise only at the second level. The necessary consistency conditions for (4.29) then follow from the requirement

$$\{Q^i{}_\alpha, S_l^\gamma\}Q^i{}_\beta|R,r\rangle^{\rm hw}_{\dot\alpha_1\ldots\dot\alpha_{2\bar\jmath}}=Q^i{}_\alpha\{Q^i{}_\beta,S_l^\gamma\}|R,r\rangle^{\rm hw}_{\dot\alpha_1\ldots\dot\alpha_{2\bar\jmath}}\,,\qquad l=1,2\,. \tag{4.30}$$

Using (3.4a) with (3.8), (3.7) and (3.11) then gives

$$\big(\delta^i{}_l(-1+\tfrac{1}{2}\Delta)-R^i{}_l\big)|R,r\rangle^{\rm hw}_{\dot\alpha_1\ldots\dot\alpha_{2\bar\jmath}}=0\,,\qquad l=1,2\,. \tag{4.31}$$

Taking (4.29) to be valid for just $i=1$ we obtain

$$\Delta=2+2R+r\,, \tag{4.32}$$

and, instead of (4.27), the action of the Q supercharges on this highest weight state gives

$$R_{(0,\bar\jmath)} \xrightarrow{Q} (R\pm\tfrac{1}{2})_{(\frac{1}{2},\bar\jmath)} \xrightarrow{Q^2} \begin{matrix} R_{(1,\bar\jmath),(0,\bar\jmath)} \\ (R-1)_{(0,\bar\jmath)}\end{matrix} \xrightarrow{Q^3} (R-\tfrac{1}{2})_{(\frac{1}{2},\bar\jmath)}\,. \tag{4.33}$$

For both $i=1,2$ (4.29) requires $R=0$ and $\Delta=2+r$. Note also that $(Q^1)^2|0,r\rangle=0$ also entails $\varepsilon^{\alpha\beta}Q^1{}_\alpha Q^2{}_\beta|0,r\rangle=(Q^2)^2|0,r\rangle=0$.

If the semi-shortening conditions (4.24) or (4.29) are applied to a superconformal primary state, with scale dimensions given by (4.26) or (4.32), then together with the action of the $\bar{Q}$ supercharges, given by the conjugate of (4.8), leads to a supermultiplet $\mathcal{C}_{R,r(j,\bar\jmath)}$ with

$$\dim\mathcal{C}_{R,r(j,\bar\jmath)}=128(R(4j+3)+j+1)(2\bar\jmath+1)\,. \tag{4.34}$$

In a similar fashion we may impose the conjugate conditions to (4.24) or (4.29),

$$\begin{aligned}\varepsilon^{\dot\alpha\dot\beta}\bar{Q}_{i\dot\beta}|R,r\rangle^{\rm hw}_{\alpha_1\ldots\alpha_{2j},\dot\alpha_1\ldots\dot\alpha_{2\bar\jmath-1}\dot\alpha}=0\,,\quad \bar\jmath>0\,,\\ (\bar{Q}_i)^2|R,r\rangle^{\rm hw}_{\alpha_1\ldots\alpha_{2j}}=0\,,\quad \bar\jmath=0\,,\end{aligned} \tag{4.35}$$

giving for just $i = 2$

$$\Delta = 2 + 2\bar{\jmath} + 2R - r\,, \qquad \Delta = 2 + 2R - r\,, \tag{4.36}$$

in each case. The action of $\bar{Q}$ is then given by the conjugate of (4.27) or (4.33) respectively. Requiring (4.35) for $i = 1, 2$ is identical as above after taking $R = 0$. Assuming (4.35) for $i = 2$ leads to a supermultiplet $\bar{\mathcal{C}}_{R,r(j,\bar{\jmath})}$ which is the conjugate of $\mathcal{C}_{R,r(j,\bar{\jmath})}$.

If the semi-shortening conditions are applied for both the Q and $\bar{Q}$ supercharges then a semi-short supermultiplet denoted by $\hat{\mathcal{C}}_{R(j,\bar{\jmath})}$ is obtained. Assuming both (4.24) and (4.35) then for compatibility of (4.26) and (4.36) $r = \bar{\jmath} - j$. Just as in in other cases we may easily construct a diagram displaying all representations that appear in the supermultiplet in this case by using (4.27) and its conjugate to obtain those formed by the action of both Q's and $\bar{Q}$'s,



(4.37)

For this case the dimension formula is

$$\dim \hat{\mathcal{C}}_{R(j,\bar{\jmath})} = 32R(4j+3)(4\bar{\jmath}+3) + 32(j+\bar{\jmath}+\tfrac{3}{2})\,. \tag{4.38}$$

For $j=\bar{\jmath}=0$ there is no restriction on the representations appearing at the first level.

If $R=0$ we may impose both the conditions (4.24) and (4.35) simultaneously for $i=1,2$. In this case compatibility with (3.1) requires that we have

$$\tilde{\mathrm{P}}^{\dot{\alpha}\alpha}|0,\bar{\jmath}{-}j\rangle_{\alpha\alpha_1\ldots\alpha_{2j-1},\dot{\alpha}\dot{\alpha}_1\ldots\dot{\alpha}_{2\bar{\jmath}-1}} = 0\,, \tag{4.39}$$

where $\Delta = 2+j+\bar{\jmath}$ for this state. This is of course a generalised conservation equation. The full multiplet $\hat{\mathcal{C}}_{0(j,\bar{\jmath})}$ is then described by



(4.40)

Every state satisfies a generalised conservation equation akin to (4.39). Taking into account these constraints the dimension is still given by (4.38) for $R=0$. If $j=0$ then it is appropriate to impose (4.29) and (4.35) for $i=1,2$. In this case (3.1) leads to

$$\tilde{\mathrm{P}}^{\dot{\alpha}\alpha}Q^i{}_\alpha|0,\bar{\jmath}\rangle_{\dot{\alpha}\dot{\alpha}_1\ldots\dot{\alpha}_{2\bar{\jmath}-1}} = 0\,, \tag{4.41}$$

which leads to the conservation equation holding for all representations $R_(j,\bar{\jmath})$ in the multiplet with $j,\bar{\jmath}>0$. A similar result of course holds if $\bar{\jmath}=0$. If $j=\bar{\jmath}=0$ then applying (4.29) and (4.35) for $i=1,2$ gives

$$\tilde{\mathrm{P}}^{\dot{\alpha}\alpha}[Q^i{}_\alpha, \bar{Q}_{j\dot{\alpha}}]|0,0\rangle = 0\,, \tag{4.42}$$

which implies that the conservation equations apply first at the second level. This multiplet, in which the lowest dimension state has $r = 0$ and $\Delta = 2$, contains the conserved currents for $U(2)_R$ symmetry as well as the conserved energy momentum tensor and was first constructed by Sohnius [21]. Counting states gives a dimension of 24 in accord with (4.38).

At the unitarity threshold given by (4.18) we may apply (4.35) to reduce further the supermultiplet displayed in (4.16) giving an asymmetric multiplet which we denote by $\mathcal{D}_{R(0,\bar{\jmath})}$,



(4.43)

In this case

$$\dim \mathcal{D}_{R(0,\bar{\jmath})} = 16(4R-1)(2\bar{\jmath}+1) + 32R\,. \tag{4.44}$$

We may clearly also define a conjugate multiplet $\bar{\mathcal{D}}_{R(j,0)}$ if $-r = j+1$ whose dimension is the same as (4.44) with $\bar{\jmath} \to j$.

For chiral multiplets, where the lowest dimension states have $R = 0$ and satisfy $Q^i{}_\alpha|0,r\rangle_{\dot\alpha_1\ldots\dot\alpha_{2\bar\jmath}} = 0$ or $\bar{Q}_{i\dot\alpha}|0,r\rangle_{\alpha_1\ldots\alpha_{2j}} = 0$ for $i = 1,2$, we may also impose the semi-shortening conditions (4.35), for $\Delta = r = 1+\bar{\jmath}$, or (4.24), for $\Delta = -r = 1+j$ respectively. For compatibility with (3.1) these states must satisfy a generalised Dirac equation, so that in the latter case we have

$$\tilde{\mathrm{P}}^{\dot\alpha\alpha}|0,-1-j\rangle_{\alpha\alpha_1\ldots\alpha_{2j-1}} = 0\,. \tag{4.45}$$

Taking $j = \bar{j}$ the reduced chiral multiplet and its conjugate are represented by



(4.46)

For a $(j, 0)$ or $(0, \bar{j})$ field obeying the Dirac equation there is one degree of freedom so that each multiplet in (4.46) has dimension 4. If $j = 0$ the lowest state satisfies $P^2|0, \pm 1\rangle = 0$ although at higher levels the states satisfy the generalised Dirac equation. This describes the standard $\mathcal{N} = 2$ vector supermultiplet, whose lowest dimension state is a massless scalar with scale dimension 1.

The semi-short supermultiplet $\hat{\mathcal{C}}_{R(j,\bar{j})}$ represented by (4.37) occurs at the threshold of the unitarity bound (4.10). It is interesting to consider how a long supermultiplet $\mathcal{A}^{\Delta}_{R,r(j,\bar{j})}$ decomposes at this point. For $j, \bar{j} > 0$ it can be written as a semi-direct sum of semi-short representations,

$$\mathcal{A}^{2R+j+\bar{j}+2}_{R,\bar{j}-j(j,\bar{j})} \simeq \hat{\mathcal{C}}_{R(j,\bar{j})} \oplus \hat{\mathcal{C}}_{R+\frac{1}{2}(j-\frac{1}{2},\bar{j})} \oplus \hat{\mathcal{C}}_{R+\frac{1}{2}(j,\bar{j}-\frac{1}{2})} \oplus \hat{\mathcal{C}}_{R+1(j-\frac{1}{2},\bar{j}-\frac{1}{2})} \,. \tag{4.47}$$

For $\bar{j} = 0$ we have

$$\mathcal{A}^{2R+j+2}_{R,-j(j,0)} \simeq \hat{\mathcal{C}}_{R(j,0)} \oplus \hat{\mathcal{C}}_{R+\frac{1}{2}(j-\frac{1}{2},0)} \oplus \bar{\mathcal{D}}_{R+1(j,0)} \oplus \bar{\mathcal{D}}_{R+\frac{3}{2}(j-\frac{1}{2},0)} \,, \tag{4.48}$$

and for $j = \bar{j} = 0$ the decomposition of $\mathcal{A}$ becomes

$$\mathcal{A}^{2R+2}_{R,0(0,0)} \simeq \hat{\mathcal{C}}_{R(0,0)} \oplus \mathcal{D}_{R+1(0,0)} \oplus \bar{\mathcal{D}}_{R+1(0,0)} \oplus \hat{\mathcal{B}}_{R+2} \,, \tag{4.49}$$

which involves a short BPS representation. We also have

$$\mathcal{B}_{R,\bar{j}+1(0,\bar{j})} \simeq \mathcal{D}_{R(0,\bar{j})} \oplus \mathcal{D}_{R+\frac{1}{2}(0,\bar{j}-\frac{1}{2})} \,, \qquad \mathcal{B}_{R,1(0,0)} \simeq \mathcal{D}_{R(0,0)} \oplus \hat{\mathcal{B}}_{R+1} \,, \tag{4.50}$$

together with their conjugates. In each case the last term represents an invariant subspace for the long multiplet. These decompositions are consistent with (4.9), (4.38), (4.17), (4.19) and (4.23). Both (4.47) and (4.49) are valid for $R = 0$ when the lowest dimension multiplet contains conserved currents. We note that (4.48) and (4.49) are special cases of (4.47) with the use of

$$\hat{\mathcal{C}}_{R(j,-\frac{1}{2})} \simeq \bar{\mathcal{D}}_{R+\frac{1}{2}(j,0)} \,, \qquad \hat{\mathcal{C}}_{R(-\frac{1}{2},\bar{j})} \simeq \mathcal{D}_{R+\frac{1}{2}(0,\bar{j})} \,, \qquad \bar{\mathcal{D}}_{R(-\frac{1}{2},0)} \simeq \mathcal{D}_{R(0,-\frac{1}{2})} \simeq \hat{\mathcal{B}}_{R+\frac{1}{2}} \,. \tag{4.51}$$

If the semi-shortening condition is applied to just the action of the Q or $\bar{Q}$ supercharges, so that only one of the bounds in (4.10) is saturated, then instead of (4.47) we have

$$\mathcal{A}^{2R+r+2j+2}_{R,r(j,\bar{\jmath})} \simeq \mathcal{C}_{R,r(j,\bar{\jmath})} \oplus \mathcal{C}_{R+\frac{1}{2},r+\frac{1}{2}(j-\frac{1}{2},\bar{\jmath})}\,, \tag{4.52}$$

together with its conjugate. Note that $\mathcal{C}_{R,r(-\frac{1}{2},\bar{\jmath})} = \mathcal{B}_{R+\frac{1}{2},r(0,\bar{\jmath})}$.

5. Superconformal Representations, $\mathcal{N} = 4$

For the $\mathcal{N} = 4$ case the discussion of possible supermultiplets is essentially similar to $\mathcal{N} = 2$. Instead of a spinorial basis as in (4.2) we now write

$$\left[M_\alpha{}^\beta\right] = \begin{pmatrix} J_3 & J_+ \\ J_- & -J_3 \end{pmatrix}\,, \qquad \left[\bar{M}^{\dot\beta}{}_{\dot\alpha}\right] = \begin{pmatrix} \bar{J}_3 & \bar{J}_+ \\ \bar{J}_- & -\bar{J}_3 \end{pmatrix}\,, \tag{5.1}$$

which satisfy (3.6) subject to the usual commutation relations for the $J_\pm$, J_3 and $\bar{J}_\pm$, $\bar{J}_3$ generators of $SU(2)_J$ and $SU(2)_{\bar{J}}$. The standard basis states are then given by $|k,p,q;m,\bar{m}\rangle$, labelled by the eigenvalues k,p,q of H_1,H_2,H_3 for $SU(4)_R$ and $m,\bar{m}$ of $J_3,\bar{J}_3$. The highest weight states then satisfy

$$E_i{}^+|k,p,q;j,\bar{\jmath}\rangle^{\rm hw} = J_+|k,p,q;j,\bar{\jmath}\rangle^{\rm hw} = \bar{J}_+|k,p,q;j,\bar{\jmath}\rangle^{\rm hw} = 0\,, \tag{5.2}$$

as well as the analogues of (4.4) and (4.5) for this to be the lowest dimension conformal primary state of a supermultiplet. The representation defined by (5.2) is here denoted by $[k,p,q]_{(j,\bar{\jmath})}$.

The supermultiplet is generated by the action of the supercharges $Q^i{}_\alpha$, $\bar{Q}_{i\dot\alpha}$ on the state with lowest scale dimension Δ. The supercharges have weights, corresponding to the change in the eigenvalues of H_i and $J_3,\bar{J}_3$, which are, from (3.27) and (3.28),

$$\begin{aligned} &Q^1{}_\alpha \sim [1,0,0]_{(\pm\frac{1}{2},0)}\,,\ Q^2{}_\alpha \sim [-1,1,0]_{(\pm\frac{1}{2},0)}\,,\ Q^3{}_\alpha \sim [0,-1,1]_{(\pm\frac{1}{2},0)}\,,\ Q^4{}_\alpha \sim [0,0,-1]_{(\pm\frac{1}{2},0)}\,,\\ &\bar{Q}_{1\dot\alpha} \sim [-1,0,0]_{(0,\pm\frac{1}{2})}\,,\ \bar{Q}_{2\dot\alpha} \sim [1,-1,0]_{(0,\pm\frac{1}{2})}\,,\ \bar{Q}_{3\dot\alpha} \sim [0,1,-1]_{(0,\pm\frac{1}{2})}\,,\ \bar{Q}_{4\dot\alpha} \sim [0,0,1]_{(0,\pm\frac{1}{2})}\,, \end{aligned} \tag{5.3}$$

which define the weights of all the Q and $\bar{Q}$ supercharges so that $Q^1{}_1$ and $\bar{Q}_{4\,1}$ are the highest weight operators of the $[1,0,0]_{(\frac{1}{2},0)}$ and $[0,0,1]_{(0,\frac{1}{2})}$ representations of $SU(4)_R \otimes SU(2)_J \otimes SU(2)_{\bar{J}}$. A long supermultiplet is generated by the unconstrained action of the 16 supercharges on the lowest dimension state, in a similar fashion to (4.6). For the Q supercharges acting on a general multiplet $[k,p,q]_{(j,\bar{\jmath})}$ we have according to the Racah-Speiser algorithm generically all possible $[k',p',q']_{(j',\bar{\jmath}')}$ obtained by adding the weights in (5.3) for each supercharge. The action of the supercharges then leads to all representations

with $[k',p',q']_{(j',\bar{j}')}$ as the highest weight, so long as $k',p',q',j',\bar{j}'$ are all positive or zero. If any of $k',p',q',j',\bar{j}'$ become negative the appropriate prescription is given in the next section. In general the representations which appear are described by

$$
\begin{aligned}
[k,p,q]_{(j,\bar{j})} \xrightarrow{Q} & \; [k{+}1,p,q],[k,p{-}1,q{+}1]_{(j\pm\frac{1}{2},\bar{j})} \\
& \; [k,p,q{-}1],[k{-}1,p{+}1,q]_{(j\pm\frac{1}{2},\bar{j})} \\
\xrightarrow{Q^2} & \; [k{+}1,p,q{-}1],[k{+}1,p{-}1,q{+}1],[k,p{+}1,q],[k,p{-}1,q],[k{-}1,p{+}1,q{-}1],[k{-}1,p,q{+}1]_{(j\pm1,\bar{j})} \\
& \qquad [k{+}2,p,q],2[k{+}1,p,q{-}1],2[k{+}1,p{-}1,q{+}1],2[k,p{+}1,q],2[k,p{-}1,q]_{(j,\bar{j})} \\
& \qquad 2[k{-}1,p{+}1,q{-}1],2[k{-}1,p,q{+}1],[k,p,q{-}2],[k,p{-}2,q{+}2],[k{-}2,p{+}2,q]_{(j,\bar{j})} \\
\xrightarrow{Q^3} & \qquad\qquad [k{+}1,p{-}1,q],[k,p{+}1,q{-}1],[k,p,q{+}1],[k{-}1,p,q]_{(j\pm\frac{3}{2},\bar{j})} \\
& \qquad [k{+}2,p,q{-}1],[k{+}2,p{-}1,q{+}1],[k{+}1,p{+}1,q],[k{+}1,p,q{-}2],[k{+}1,p{-}2,q{+}2]_{(j\pm\frac{1}{2},\bar{j})} \\
& 3[k{+}1,p{-}1,q],3[k,p{+}1,q{-}1],3[k,p,q{+}1],3[k{-}1,p,q],[k,p{-}1,q{-}1],[k,p{-}2,q{+}1]_{(j\pm\frac{1}{2},\bar{j})} \\
& [k{-}1,p{+}2,q],[k{-}1,p{+}1,q{-}2],[k{-}1,p{-}1,q{+}2],[k{-}2,p{+}2,q{-}1],[k{-}2,p{+}1,q{+}1]_{(j\pm\frac{1}{2},\bar{j})} \\
\xrightarrow{Q^4} & \; [k,p,q]_{(j\pm2,\bar{j})},[k{+}2,p{-}1,q{+}1],[k{+}1,p{+}1,q{-}1],[k{+}1,p,q{+}1],[k{+}1,p{-}1,q{+}1]_{(j\pm1,\bar{j})} \\
& \qquad [k{+}1,p{-}2,q{+}1],[k,p{+}1,q{-}2],4[k,p,q],[k,p{-}1,q{+}2],[k{-}1,p{+}1,q{+}1]_{(j\pm1,\bar{j})} \\
& \qquad\qquad [k{-}1,p,q{-}1],[k{-}1,p{-}1,q{+}1],[k{-}1,p{+}2,q{+}1],[k{-}2,p{+}1,q]_{(j\pm1,\bar{j})} \\
& \qquad\qquad\qquad [k{+}2,p,q{-}2],2[k{+}2,p{-}1,q],[k{+}2,p{-}2,q{+}2]_{(j,\bar{j})} \\
& \qquad\qquad 2[k{+}1,p{+}1,q{-}1],2[k{+}1,p,q{+}1],2[k{+}1,p{-}1,q{-}1],2[k{+}1,p{-}2,q{+}1]_{(j,\bar{j})} \\
& \qquad\qquad\qquad [k,p{+}2,q],2[k,p{+}1,q{-}2],6[k,p,q],2[k,p{-}1,q{+}2],[k,p{-}2,q]_{(j,\bar{j})} \\
& \qquad\qquad 2[k{-}1,p{+}2,q{-}1],2[k{-}1,p{+}1,q{+}1],2[k{-}1,p,q{-}1],2[k{-}1,p{-}1,q{+}1]_{(j,\bar{j})} \\
& \qquad\qquad\qquad [k{-}2,p{+}2,q{-}2],2[k{-}2,p{+}1,q],[k{-}2,p,q{+}2]_{(j,\bar{j})} \\
\xrightarrow{Q^5} & \qquad\qquad [k{+}1,p,q],[k,p,q{-}1],[k,p{-}1,q{+}1],[k{-}1,p{+}1,q]_{(j\pm\frac{3}{2},\bar{j})} \\
& \; [k{+}2,p{-}1,q{-}1],[k{+}2,p{-}2,q{+}1],[k{+}1,p{+}1,q{-}2],[k{+}1,p{-}1,q{+}2],[k{+}1,p{-}2,q]_{(j\pm\frac{1}{2},\bar{j})} \\
& 3[k{+}1,p,q],3[k,p,q{-}1],3[k,p{-}1,q{+}1],3[k{-}1,p{+}1,q],[k,p{+}2,q{-}1],[k,p{+}1,q{+}1]_{(j\pm\frac{1}{2},\bar{j})} \\
& \qquad [k{-}1,p{+}2,q{-}2],[k{-}1,p,q{+}2],[k{-}1,p{-}1,q],[k{-}2,p{+}1,q{-}1],[k{-}2,p,q{+}1]_{(j\pm\frac{1}{2},\bar{j})} \\
\xrightarrow{Q^6} & \; [k{+}1,p,q{-}1],[k{+}1,p{-}1,q{+}1],[k,p{+}1,q],[k,p{-}1,q],[k{-}1,p{+}1,q{-}1],[k{-}1,p,q{+}1]_{(j\pm1,\bar{j})} \\
& \qquad [k{+}2,p{-}2,q],2[k{+}1,p,q{-}1],2[k{+}1,p{-}1,q{+}1],2[k,p{+}1,q],2[k,p{-}1,q]_{(j,\bar{j})} \\
& \qquad 2[k{-}1,p{+}1,q{-}1],2[k{-}1,p,q{+}1],[k,p{+}2,q{-}2],[k,p,q{+}2],[k{-}2,p,q]_{(j,\bar{j})} \\
\xrightarrow{Q^7} & \; [k{+}1,p{-}1,q],[k,p{+}1,q{-}1]_{(j\pm\frac{1}{2},\bar{j})} \\
& \qquad [k,p,q{+}1],[k{-}1,p,q]_{(j\pm\frac{1}{2},\bar{j})} \\
\xrightarrow{Q^8} & \; [k,p,q]_{(j,\bar{j})} \,.
\end{aligned}
\tag{5.4}
$$

At each level l there are $\binom{8}{l}$ representations of $SU(4)_R \otimes SU(2)_J \otimes SU(2)_{\bar{J}}$. The total dimension of this module is 2^8 times the dimension of the lowest dimension representation. Defining $\hat{k} = k+1$, $\hat{p} = p+1$, $\hat{q} = q+1$ and $\mathcal{J} = 2j+1$, $\bar{\mathcal{J}} = 2\bar{j}+1$ this becomes, with the $SU(4)$ dimension formula given in (3.29),

$$
2^8 d(k,p,q)(2j+1)(2\bar{j}+1) = \frac{64}{3}\,\hat{k}\,\hat{p}\,\hat{q}(\hat{k}+\hat{p})(\hat{q}+\hat{p})(\hat{k}+\hat{p}+\hat{q})\mathcal{J}\,\bar{\mathcal{J}}\,. \tag{5.5}
$$

For multiplet shortening we consider BPS conditions of the form

$$
Q^i{}_\alpha |k,p,q;j,\bar{j}\rangle^{\rm hw} = 0\,, \qquad \alpha = 1,2\,, \tag{5.6}
$$

which restricts the representation content at level one and subsequent levels. Since, for the lowest scale dimension state, $|k,p,q;j,\bar{j}\rangle^{\rm hw}$ is annihilated by $S_l{}^\beta$ then applying (3.4a) requires that the state must also be annihilated by $M_\alpha{}^\beta$ or $j=0$ and then

$$\big(\delta^i_{\ l}\tfrac{1}{2}\Delta - R^i_{\ l}\big)|k,p,q;0,\bar{j}\rangle^{\rm hw} = 0\,, \qquad l=1,2,3,4\,. \tag{5.7}$$

Using (3.26) and (3.22), (3.23), (3.24) it is easy to see that there are the following solutions, where s is the fraction of the charges for which (5.6) holds,

$$\begin{aligned}
&i=1 && s=\tfrac{1}{4} && \Delta=\tfrac{1}{2}(3k+2p+q)\,, && \\
&i=1,2 && s=\tfrac{1}{2} && \Delta=\tfrac{1}{2}(2p+q)\,, && k=0\,,\\
&i=1,2,3 && s=\tfrac{3}{4} && \Delta=\tfrac{1}{2}q\,, && k=q=0\,,\\
&i=1,2,3,4 && s=1 && \Delta=0\,, && k=p=q=0\,.
\end{aligned} \tag{5.8}$$

The last case of course corresponds to the trivial singlet representation. Conversely since from (3.28) $[E_1{}^-,Q^1{}_\alpha]=Q^2{}_\alpha$ if $k=0$ the condition (5.6) for $i=1$ implies the $i=2$ case by virtue of (3.24). Similarly if $p=0$ or $p=q=0$ as well as $k=0$ we have the BPS condition for $i=2$ or $i=2,3$.

For each case we may obtain the set of representations, analogous to (5.4), obtained by the action of the supercharges on a lowest dimension state $[k,p,q]_{(0,\bar{j})}$. For the $\frac{1}{4}$-BPS case the six supercharges $Q^i{}_\alpha$, $i=2,3,4$, give, for general k,p,q,

$$\begin{aligned}
[k,p,q]_{(0,\bar{j})} \xrightarrow{Q}\ & [k,p,q-1],[k,p-1,q+1]_{(\frac{1}{2},\bar{j})}\\
& \quad [k-1,p+1,q]_{(\frac{1}{2},\bar{j})}\\
\xrightarrow{Q^2}\ & \quad [k,p-1,q],[k-1,p+1,q-1],[k-1,p,q+1]_{(1,\bar{j})}\\
& [k,p,q-2],[k,p-1,q],[k,p-2,q+2],[k-1,p+1,q-1],[k-1,p,q+1],[k-2,p+2,q]_{(0,\bar{j})}\\
\xrightarrow{Q^3}\ & \quad [k-1,p,q]_{(\frac{3}{2},\bar{j})},[k,p-1,q-1],[k,p-2,q+1]_{(\frac{1}{2},\bar{j})}\\
& [k-1,p+1,q-2],2[k-1,p,q],[k-1,p-1,q+2],[k-2,p+2,q-1],[k-2,p+1,q+1]_{(\frac{1}{2},\bar{j})}\\
\xrightarrow{Q^4}\ & \quad [k-1,p,q-1],[k-1,p-1,q+1],[k-2,p+1,q]_{(1,\bar{j})}\\
& [k,p-2,q],[k-1,p,q-1],[k-1,p-1,q+1],[k-2,p+2,q-2],[k-2,p+1,q],[k-2,p,q+2]_{(0,\bar{j})}\\
\xrightarrow{Q^5}\ & \quad [k-1,p-1,q]_{(\frac{1}{2},\bar{j})}\\
& [k-2,p+1,q-1],[k-2,p,q+1]_{(\frac{1}{2},\bar{j})}\\
\xrightarrow{Q^6}\ & [k-2,p,q]_{(0,\bar{j})}\,.
\end{aligned} \tag{5.9}$$

In this case at level l there are $\binom{6}{l}$ representations of $SU(4)_R\otimes SU(2)_{\bar{J}}$. The dimension of this module, which may be obtained by adding the dimensions of all representations in (5.9), is,

$$2^6 d(k-1,p,q)\bar{\mathcal{J}} - \tfrac{8}{3}\,\hat{p}\,\hat{q}(\hat{p}+\hat{q})(3\hat{k}+2\hat{p}+\hat{q}-3)\bar{\mathcal{J}}\,. \tag{5.10}$$

When the $\frac{1}{2}$-BPS condition is applied the states are obtained just by the action of the ${Q^3}_\alpha, {Q^4}_\alpha$ supercharges. According to (5.8) we should set $k=0$ in the lowest dimension state for this case but we allow for arbitrary k for subsequent appplication. The action of the supercharges then gives representations obtained by adding weights,

$$\begin{aligned}[k,p,q]_{(0,\bar{j})} \xrightarrow{Q} & \begin{matrix}[k,p,q-1]_{(\frac{1}{2},\bar{j})}\\ [k,p-1,q+1]_{(\frac{1}{2},\bar{j})}\end{matrix} \xrightarrow{Q^2} \begin{matrix}[k,p-1,q]_{(1,\bar{j})},[k,p,q-2]_{(0,\bar{j})}\\ [k,p-1,q],[k,p-2,q+2]_{(0,\bar{j})}\end{matrix}\\ \xrightarrow{Q^3} & \begin{matrix}[k,p-1,q-1]_{(\frac{1}{2},\bar{j})}\\ [k,p-2,q+1]_{(\frac{1}{2},\bar{j})}\end{matrix} \xrightarrow{Q^4} [k,p-2,q]_{(0,\bar{j})}\,. \end{aligned} \tag{5.11}$$

In (5.11) it is easy to see that there are now $\binom{4}{l}$ representations of $SU(4)_R \otimes SU(2)_{\bar{J}}$. The result (5.11) is valid for $k \geq 0$, $p,q \geq 2$, restriction to other cases will be given later. The dimension for module given by (5.11) is,

$$2^4 d(k,p-1,q)\bar{\mathcal{J}} - \tfrac{1}{3}\hat{k}\,\hat{q}(2\hat{k}^2 - 2\hat{q}^2 - 3)\bar{\mathcal{J}}\,. \tag{5.12}$$

For completeness we also note the $\frac{3}{4}$-BPS case which is obtained by the action of ${Q^4}_\alpha$ alone

$$[k,p,q]_{(0,\bar{j})} \xrightarrow{Q} [k,p,q-1]_{(\frac{1}{2},\bar{j})} \xrightarrow{Q^2} [k,p,q-2]_{(0,\bar{j})}\,, \tag{5.13}$$

again generalising to arbitrary k,p,q, with dimension

$$2^2 d(k,p,q-1)\bar{\mathcal{J}} + \tfrac{1}{6}\,\hat{k}\,\hat{p}(\hat{k}+\hat{p})(\hat{k}+2\hat{p}+3\hat{q}-3)\bar{\mathcal{J}}\,. \tag{5.14}$$

As in the $\mathcal{N}=2$ case there are further shortening conditions that may be applied for $j \geq 0$. We consider the constraint

$$\Big(Q^i{}_2 - \tfrac{1}{2j+1} J_- Q^i{}_1\Big)|k,p,q;j,\bar{j}\rangle^{\rm hw} = 0\,. \tag{5.15}$$

Since J_+ acting on this state gives zero, using $[J_+, Q^i{}_2] = Q^i{}_1$, this is clearly a state with spin $(j-1,\bar{j})$ and hence (5.15) is equivalent for $\mathcal{N}=4$ to (4.24) for $\mathcal{N}=2$. Applying S^1_l to this and using $\{S^1_l, Q^i{}_2\} = 4\delta^i{}_l J_-$, $[S^1_l, J_-] = 0$, $\{S^1_l, Q^i{}_1\} = 4\delta^i{}_l(J_3 - \frac{1}{2}iD) + 4R^i{}_l$, the condition (5.15) leads to the constraint

$$J_-\Big(\delta^i{}_l - \tfrac{1}{2j+1}\big((j+\tfrac{1}{2}\Delta)\delta^i{}_l - R^i{}_l\big)\Big)|k,p,q;j,\bar{j}\rangle^{\rm hw} = 0\,, \quad l=1,2,3,4\,, \tag{5.16}$$

which determines Δ. Since $S^2_l = -[J_+, S^1_l]$ it is clear that applying S^2_l to (5.15) does not give additional conditions. Again there are various cases according to the various possible choices i for which (5.15) holds,

$$\begin{aligned}
&i=1 && t=\tfrac{1}{4} && \Delta = 2+2j+\tfrac{1}{2}(3k+2p+q)\,,\\
&i=1,2 && t=\tfrac{1}{2} && \Delta = 2+2j+\tfrac{1}{2}(2p+q)\,, && k=0\,,\\
&i=1,2,3 && t=\tfrac{3}{4} && \Delta = 2+2j+\tfrac{1}{2}q\,, && k=p=0\,,\\
&i=1,2,3,4 && t=1 && \Delta = 2+2j\,, && k=p=q=0\,,
\end{aligned} \tag{5.17}$$

where t is analogous to s in (5.8). As discussed earlier $p = 0$ requires that (5.15) for $i = 1$ implies $i = 2$, $k = p = 0$ implies (5.15) for $i = 2, 3$ and $k = p = q = 0$ for $i = 2, 3, 4$.

Applying the condition (5.15) just for $i = 1$ leads to the module

$$
\begin{aligned}
[k,p,q]_{(j,\bar{\jmath})} \xrightarrow{Q} & \; [k{+}1,p,q]_{(j+\frac{1}{2},\bar{\jmath})},[k,p{-}1,q{+}1]_{(j\pm\frac{1}{2},\bar{\jmath})} \\
& \quad [k,p,q{-}1],[k{-}1,p{+}1,q]_{(j\pm\frac{1}{2},\bar{\jmath})} \\
\xrightarrow{Q^2} & \; [k{+}1,p,q{-}1],[k{+}1,p{-}1,q{+}1],[k,p{+}1,q]_{(j+1,\bar{\jmath})},[k,p{-}1,q],[k{-}1,p{+}1,q{-}1],[k{-}1,p,q{+}1]_{(j\pm1,\bar{\jmath})} \\
& \quad [k{+}1,p,q{-}1],[k{+}1,p{-}1,q{+}1],[k,p{+}1,q],2[k,p{-}1,q]_{(j,\bar{\jmath})} \\
& \quad 2[k{-}1,p{+}1,q{-}1],2[k{-}1,p,q{+}1],[k,p,q{-}2],[k,p{-}2,q{+}2],[k{-}2,p{+}2,q]_{(j,\bar{\jmath})} \\
\xrightarrow{Q^3} & \; [k{+}1,p{-}1,q],[k,p{+}1,q{-}1],[k,p,q{+}1]_{(j+\frac{3}{2},\bar{\jmath})},[k{-}1,p,q]_{(j\pm\frac{3}{2},\bar{\jmath})} \\
& \quad [k{+}1,p,q{-}2],[k{+}1,p{-}1,q],[k{+}1,p{-}2,q{+}2],[k,p{+}1,q{-}1],[k,p,q{+}1],k{-}1,p{+}2,q]_{(j+\frac{1}{2},\bar{\jmath})} \\
& \quad [k{+}1,p{-}1,q],[k,p{+}1,q{-}1],[k,p,q{+}1],[k,p{-}1,q{-}1],[k,p{-}2,q{+}1]_{(j\pm\frac{1}{2},\bar{\jmath})} \\
& \quad [k{-}1,p{+}1,q{-}2],3[k{-}1,p,q],[k{-}1,p{-}1,q{+}2],[k{-}2,p{+}2,q{-}1],[k{-}2,p{+}1,q{+}1]_{(j\pm\frac{1}{2},\bar{\jmath})} \\
\xrightarrow{Q^4} & \; [k,p,q]_{(j+2,\bar{\jmath})},[k{+}1,p{-}1,q{-}1],[k{+}1,p{-}2,q{+}1],[k,p{+}1,q{-}2],2[k,p,q]_{(j+1,\bar{\jmath})} \\
& \quad [k,p{-}1,q{+}2],[k{-}1,p{+}2,q{-}1],[k{-}1,p{+}1,q{+}1]_{(j+1,\bar{\jmath})} \\
& \quad [k,p,q],[k{-}1,p,q{-}1],[k{-}1,p{-}1,q{+}1],[k{-}2,p{+}1,q]_{(j\pm1,\bar{\jmath})} \\
& \quad [k{+}1,p{-}1,q{-}1],[k{+}1,p{-}2,q{+}1],[k,p{+}1,q{-}2],3[k,p,q],[k,p{-}1,q{+}2],[k,p{-}2,q]_{(j,\bar{\jmath})} \\
& \quad [k{-}1,p{+}2,q{-}1],[k{-}1,p{+}1,q{+}1],2[k{-}1,p,q{-}1],2[k{-}1,p{-}1,q{+}1]_{(j,\bar{\jmath})} \\
& \quad [k{-}2,p{+}2,q{-}2],2[k{-}2,p{+}1,q],[k{-}2,p,q{+}2]_{(j,\bar{\jmath})} \\
\xrightarrow{Q^5} & \; [k,p,q{-}1],[k,p{-}1,q{+}1],[k{-}1,p{+}1,q]_{(j+\frac{3}{2},\bar{\jmath})} \\
& \quad [k{+}1,p{-}2,q],[k,p,q{-}1],[k,p{-}1,q{+}1],[k{-}1,p{+}2,q{-}2],[k{-}1,p{+}1,q],[k{-}1,p,q{+}2]_{(j+\frac{1}{2},\bar{\jmath})} \\
& \quad [k,p,q{-}1],[k,p{-}1,q{+}1],[k{-}1,p{+}1,q],[k{-}1,p{-}1,q],[k{-}2,p{+}1,q{-}1],[k{-}2,p,q{+}1]_{(j\pm\frac{1}{2},\bar{\jmath})} \\
\xrightarrow{Q^6} & \; [k,p{-}1,q],[k{-}1,p{+}1,q{-}1],[k{-}1,p,q{+}1]_{(j+1,\bar{\jmath})} \\
& \quad [k,p{-}1,q],[k{-}1,p{+}1,q{-}1],[k{-}1,p,q{+}1],[k{-}2,p,q]_{(j,\bar{\jmath})} \\
\xrightarrow{Q^7} & \; [k{-}1,p,q]_{(j+\frac{1}{2},\bar{\jmath})}\,.
\end{aligned}
\tag{5.18}
$$

For this case there are $\binom{7}{l}$ representations of $SU(4)_R \otimes SU(2)_J \otimes SU(2)_{\bar{J}}$ at level l. By adding up the dimensions of all representations we find a total dimension of

$$
\begin{aligned}
&\big(2^7 d(k-\tfrac{1}{2},p,q)(\mathcal{J}+\tfrac{1}{2}) - \tfrac{4}{3}\,\hat{p}\hat{q}(\hat{p}+\hat{q})(6\,\hat{k}+4\,\hat{p}+2\,\hat{q}-3)(\mathcal{J}+\tfrac{3}{2}) \\
&\quad + \tfrac{8}{3}\,\hat{p}\hat{q}(\hat{p}+\hat{q})(4\,\hat{k}\hat{p}+2\,\hat{k}\hat{q}+\hat{p}\hat{q}+\hat{p}^2+3\,\hat{k}^2-2)\big)\bar{\mathcal{J}}\,.
\end{aligned}
\tag{5.19}
$$

If (5.15) is applied for $i = 1, 2$ we obtain

$$
\begin{aligned}
[k,p,q]_{(j,\bar{\jmath})} \xrightarrow{Q} & \; [k{+}1,p,q],[k{-}1,p{+}1,q]_{(j+\frac{1}{2},\bar{\jmath})} \\
& \quad [k,p,q{-}1],[k,p{-}1,q{+}1]_{(j\pm\frac{1}{2},\bar{\jmath})} \\
\xrightarrow{Q^2} & \; [k{+}1,p,q{-}1],[k{+}1,p{-}1,q{+}1],[k,p{+}1,q],[k{-}1,p{+}1,q{-}1],[k{-}1,p,q{+}1]_{(j+1,\bar{\jmath})},[k,p{-}1,q]_{(j\pm1,\bar{\jmath})} \\
& \quad [k{+}1,p,q{-}1],[k{+}1,p{-}1,q{+}1],[k,p,q{-}2],2[k,p{-}1,q],[k,p{-}2,q{+}2],[k{-}1,p{+}1,q{-}1],[k{-}1,p,q{+}1]_{(j,\bar{\jmath})} \\
\xrightarrow{Q^3} & \; [k{+}1,p{-}1,q],[k,p{+}1,q{-}1],[k,p,q{+}1],[k{-}1,p,q]_{(j+\frac{3}{2},\bar{\jmath})} \\
& \quad [k{+}1,p,q{-}2],[k{+}1,p{-}1,q],[k{+}1,p{-}2,q{+}2],[k,p{+}1,q{-}1],[k,p,q{+}1]_{(j+\frac{1}{2},\bar{\jmath})} \\
& \quad [k{-}1,p{+}1,q{-}2],[k{-}1,p,q],[k{-}1,p{-}1,q{+}2]_{(j+\frac{1}{2},\bar{\jmath})} \\
& \quad [k{+}1,p{-}1,q],[k,p{-}1,q{-}1],[k,p{-}2,q{+}1],[k{-}1,p,q]_{(j\pm\frac{1}{2},\bar{\jmath})}
\end{aligned}
$$

$$
\begin{aligned}
&\xrightarrow{Q^4} \begin{array}{c} [k,p,q]_{(j+2,\bar{\jmath})},[k{+}1,p{-}1,q{-}1],[k{+}1,p{-}2,q{+}1],[k,p{+}1,q{-}2]_{(j+1,\bar{\jmath})} \\ 2[k,p,q],[k,p{-}1,q{+}2],[k{-}1,p,q{-}1],[k{-}1,p{-}1,q{+}1]_{(j+1,\bar{\jmath})} \\ [k{+}1,p{-}1,q{-}1],[k{+}1,p{-}2,q{+}1],[k,p,q],[k,p{-}2,q],[k{-}1,p,q{-}1],[k{-}1,p{-}1,q{+}1]_{(j,\bar{\jmath})} \end{array} \\
&\xrightarrow{Q^5} \begin{array}{c} [k,p,q{-}1],[k,p{-}1,q{+}1]_{(j+\frac{3}{2},\bar{\jmath})} \\ [k{+}1,p{-}2,q],[k,p,q{-}1],[k,p{-}1,q{+}1],[k{-}1,p{-}1,q]_{(j+\frac{1}{2},\bar{\jmath})} \end{array} \\
&\xrightarrow{Q^6} [k,p{-}1,q]_{(j+1,\bar{\jmath})}\,.
\end{aligned}
\tag{5.20}
$$

Here there are $\binom{6}{l}$ representations of $SU(4)_R \otimes SU(2)_J \otimes SU(2)_{\bar{J}}$ at level l. The total dimension of all representations is

$$
\begin{aligned}
&\big(2^6 d(k,p-\tfrac{1}{2},q)(\mathcal{J}+1) - \tfrac{1}{3}\,\hat{k}\hat{q}(4\,\hat{k}^2 - 4\,\hat{q}^2 - 3)(\mathcal{J}+1) \\
&\quad + \tfrac{4}{3}\,\hat{k}\hat{q}(\hat{k} + 2\,\hat{p} + \hat{q} - 1)(2\,\hat{k}\hat{p} + 2\,\hat{q}\hat{p} + 2\,\hat{p}^2 + \hat{k}\hat{q} - \hat{k} - 2\,\hat{p} - \hat{q} + 1)\big)\bar{\mathcal{J}}\,.
\end{aligned}
\tag{5.21}
$$

With $i = 1,2,3$ we have

$$
\begin{aligned}
[k,p,q]_{(j,\bar{\jmath})} &\xrightarrow{Q} \begin{array}{c} [k{+}1,p,q],[k,p{-}1,q{+}1],[k{-}1,p{+}1,q]_{(j+\frac{1}{2},\bar{\jmath})} \\ [k,p,q{-}1]_{(j\pm\frac{1}{2},\bar{\jmath})} \end{array} \\
&\xrightarrow{Q^2} \begin{array}{c} [k{+}1,p,q{-}1],[k{+}1,p{-}1,q{+}1],[k,p{+}1,q],[k,p{-}1,q],[k{-}1,p{+}1,q{-}1],[k{-}1,p,q{+}1]_{(j+1,\bar{\jmath})} \\ [k{+}1,p,q{-}1],[k,p,q{-}2],[k,p{-}1,q],[k{-}1,p{+}1,q{-}1]_{(j,\bar{\jmath})} \end{array} \\
&\xrightarrow{Q^3} \begin{array}{c} [k{+}1,p{-}1,q],[k,p{+}1,q{-}1],[k,p,q{+}1],[k{-}1,p,q]_{(j+\frac{3}{2},\bar{\jmath})} \\ [k{+}1,p,q{-}2],[k{+}1,p{-}1,q],[k,p{+}1,q{-}1],[k,p{-}1,q{-}1],[k{-}1,p{+}1,q{-}2],[k{-}1,p,q]_{(j+\frac{1}{2},\bar{\jmath})} \end{array} \\
&\xrightarrow{Q^4} \begin{array}{c} [k,p,q]_{(j+2,\bar{\jmath})} \\ [k{+}1,p{-}1,q{-}1],[k,p{+}1,q{-}2],[k,p,q],[k{-}1,p,q{-}1]_{(j+1,\bar{\jmath})} \end{array} \\
&\xrightarrow{Q^5} [k,p,q{-}1]_{(j+\frac{3}{2},\bar{\jmath})}\,.
\end{aligned}
\tag{5.22}
$$

The dimensions here add up to

$$
\begin{aligned}
&\big(2^5 d(k,p,q-\tfrac{1}{2})(\mathcal{J}+\tfrac{3}{2}) + \tfrac{1}{3}\,\hat{k}\hat{p}(\hat{k}+\hat{p})(2\,\hat{k} + 4\,\hat{p} + 6\,\hat{q} - 3)(\mathcal{J}+\tfrac{1}{2}) \\
&\quad + \tfrac{1}{3}\,\hat{k}\hat{p}(\hat{k}+\hat{p})(2\,\hat{k}\hat{p} + 4\,\hat{k}\hat{q} + 8\,\hat{p}\hat{q} + 2\,\hat{p}^2 + 6\,\hat{q}^2 - 1)\big)\bar{\mathcal{J}}\,.
\end{aligned}
\tag{5.23}
$$

Finally for $i = 1,2,3,4$ in (5.15),

$$
\begin{aligned}
[k,p,q]_{(j,\bar{\jmath})} &\xrightarrow{Q} \begin{array}{c} [k{+}1,p,q],[k,p,q{-}1]_{(j+\frac{1}{2},\bar{\jmath})} \\ [k,p{-}1,q{+}1,[k{-}1,p{+}1,q]_{(j+\frac{1}{2},\bar{\jmath})} \end{array} \xrightarrow{Q^2} \begin{array}{c} [k{+}1,p,q{-}1],[k{+}1,p{-}1,q{+}1],[k,p{+}1,q]_{(j+1,\bar{\jmath})} \\ [k,p{-}1,q],[k{-}1,p{+}1,q{-}1],[k{-}1,p,q{+}1]_{(j+1,\bar{\jmath})} \end{array} \\
&\xrightarrow{Q^3} \begin{array}{c} [k{+}1,p{-}1,q],[k,p{+}1,q{-}1]_{(j+\frac{3}{2},\bar{\jmath})} \\ [k,p,q{+}1],[k{-}1,p,q]_{(j+\frac{3}{2},\bar{\jmath})} \end{array} \xrightarrow{Q^4} [k,p,q]_{(j+2,\bar{\jmath})}\,,
\end{aligned}
\tag{5.24}
$$

In this case the dimension formula is

$$
2^4 d(k,p,q)(2j+3)(2\bar{\jmath}+1) = \tfrac{4}{3}\,\hat{k}\,\hat{p}\,\hat{q}(\hat{k}+\hat{p})(\hat{q}+\hat{p})(\hat{k}+\hat{p}+\hat{q})(\mathcal{J}+2)\,\bar{\mathcal{J}}\,. \tag{5.25}
$$

Equivalent shortening conditions may be obtained for the action of the $\bar{Q}$ supercharges. Instead of (5.6) we may impose

$$\bar{Q}_{i\dot{\alpha}}|k,p,q;j,\bar{\jmath}\rangle^{\rm hw}=0\,,\qquad \dot{\alpha}=1,2\,. \tag{5.26}$$

Consistency requires $\bar{\jmath}=0$ and then

$$\big(\delta^l{}_i\tfrac{1}{2}\Delta+R^l{}_i\big)|k,p,q;0,\bar{\jmath}\rangle^{\rm hw}=0\,,\qquad l=1,2,3,4\,, \tag{5.27}$$

which leads to, corresponding to (5.8),

$$\begin{array}{llll} i=4 & \bar{s}=\frac{1}{4} & \Delta=\frac{1}{2}(k+2p+3q)\,, & \\ i=3,4 & \bar{s}=\frac{1}{2} & \Delta=\frac{1}{2}(k+2p)\,, & q=0\,, \\ i=2,3,4 & \bar{s}=\frac{3}{4} & \Delta=\frac{1}{2}k\,, & p=q=0\,, \\ i=1,2,3,4 & \bar{s}=1 & \Delta=0\,, & k=p=q=0\,. \end{array} \tag{5.28}$$

For the semi-short case we may impose, corresponding to (5.15),

$$\Big(\bar{Q}_{i1}+\tfrac{1}{2\bar{\jmath}+1}\bar{J}_-\bar{Q}_{i2}\Big)|k,p,q;j,\bar{\jmath}\rangle^{\rm hw}=0\,, \tag{5.29}$$

and this leads to

$$\begin{array}{llll} i=4 & \bar{t}=\frac{1}{4} & \Delta=2+2\bar{\jmath}+\frac{1}{2}(k+2p+3q)\,, & \\ i=3,4 & \bar{t}=\frac{1}{2} & \Delta=2+2\bar{\jmath}+\frac{1}{2}(k+2p)\,, & q=0\,, \\ i=2,3,4 & \bar{t}=\frac{3}{4} & \Delta=2+2\bar{\jmath}+\frac{1}{2}k\,, & p=q=0\,, \\ i=1,2,3,4 & \bar{t}=1 & \Delta=2+2\bar{\jmath}\,, & k=p=q=0\,. \end{array} \tag{5.30}$$

For both the short and semi-short cases the action of the $\bar{Q}$ charges may be obtained by the conjugation of the results for Q given above.

The above results, summarised in (5.8) and (5.17), correspond to representations of the chiral subalgebra formed by $\{Q^i{}_\alpha, S_j{}^\beta, M_\alpha{}^\beta, R^i{}_j, D\}$. Neglecting $\bar{\jmath}$ which is irrelevant the long multiplet given by (5.4) may be denoted by

$$a^{\Delta}_{[k,p,q]j}\,, \tag{5.31}$$

while for the BPS short multiplets we have, for $j=0$,

$$b^{\frac{1}{4}}_{[k,p,q]}\,,\qquad b^{\frac{1}{2}}_{[0,p,q]}\,,\qquad b^{\frac{3}{4}}_{[0,0,q]}\,, \tag{5.32}$$

which correspond to (5.9), (5.11) and (5.13), and the semi-short multiplets are

$$c^{\frac{1}{4}}_{[k,p,q]j}\,,\qquad c^{\frac{1}{2}}_{[0,p,q]j}\,,\qquad c^{\frac{3}{4}}_{[0,0,q]j}\,,\qquad c^{1}_{[0,0,0]j}\,, \tag{5.33}$$

corresponding to (5.18) for $k \geq 1$, (5.20) for $k = 0$, $p \geq 1$, (5.22) for $k = p = 0$, $q \geq 1$ and (5.24) $k = p = q = 0$ respectively. At the unitarity threshold it is easy to then verify that the long multiplet may be decomposed as semi-direct sum

$$a^{2+2j+\frac{1}{2}(3k+2p+q)}_{[k,p,q]j} \simeq c^{\frac{1}{4}}_{[k,p,q]j} \oplus c^{\frac{1}{4}}_{[k+1,p,q]j-\frac{1}{2}}\,, \qquad a^{2+2j+\frac{1}{2}(2p+q)}_{[0,p,q]j} \simeq c^{\frac{1}{2}}_{[0,p,q]j} \oplus c^{\frac{1}{4}}_{[1,p,q]j-\frac{1}{2}}\,,$$
$$a^{2+2j+\frac{1}{2}q}_{[0,0,q]j} \simeq c^{\frac{3}{4}}_{[0,0,q]j} \oplus c^{\frac{1}{4}}_{[1,0,q]j-\frac{1}{2}}\,, \qquad a^{2+2j}_{[0,0,0]j} \simeq c^{1}_{[0,0,0]j} \oplus c^{\frac{1}{4}}_{[1,0,0]j-\frac{1}{2}}\,, \tag{5.34}$$

which can be checked with the aid of the various dimension formulae. These results also hold for $j = 0$ if we make the identification, justified later, that

$$c^{\frac{1}{4}}_{[k,p,q]-\frac{1}{2}} \simeq b^{\frac{1}{4}}_{[k+1,p,q]}\,, \tag{5.35}$$

which is consistent with the dimension results (5.19), for $\mathcal{J} = 0$, and (5.10). The first case in (5.34) has also been obtained recently in [22] and BPS multiplets further discussed in [23].

To construct all conformal primary states for complete supermultiplets it is necessary to combine the results for the Q and $\bar{Q}$ supercharges. The basic long supermultiplet is labelled in terms of its scale dimension and $SU(4)$, spin representations for the lowest dimension state,

$$\mathcal{A}^{\Delta}_{[k,p,q](j,\bar{j})}\,, \qquad \dim \mathcal{A}^{\Delta}_{[k,p,q](j,\bar{j})} = 2^{16} d(k,p,q)(2j+1)(2\bar{j}+1)\,. \tag{5.36}$$

All conformal primary representations may be found with the aid of (5.4) and its conjugate, applying this to each representation appearing in (5.4). If k, p or q are < 4 there will be a cancellation of representations according to rules specified later. For unitarity [8]

$$\Delta \geq 2 + 2j + \tfrac{1}{2}(3k + 2p + q),\ 2 + 2\bar{j} + \tfrac{1}{2}(k + 2p + 3q)\,. \tag{5.37}$$

From the above it is simple to obtain the short supermultiplets where the shortening conditions apply only to the action of the Q supercharges while $\bar{Q}$ are unconstrained, and vice versa. The full supermultiplets are easily obtained from the results above and in consequence we do not consider further such cases here.

If the conditions (5.8) and (5.28) are both applied we must have $j = \bar{j} = 0$ and we denote these short supermultiplets as $\mathcal{B}^{s,\bar{s}}_{[k,p,q](0,0)}$. It is easy to see, from compatibility of the formulae for Δ in (5.8) and (5.28), that it is necessary that $s = \bar{s}$ if they are non zero. For $s = \bar{s} = 1$, $k = p = q = 0$, we must also have from (3.1) $P_a|0,0,0;0,0\rangle = 0$ and of course this corresponds to the trivial vacuum representation. Otherwise only the two well known cases are possible,

$$\mathcal{B}^{\frac{1}{4},\frac{1}{4}}_{[q,p,q](0,0)}\,, \qquad \Delta = p + 2q\,, \tag{5.38}$$

and

$$\mathcal{B}^{\frac{1}{2},\frac{1}{2}}_{[0,p,0](0,0)}\,, \qquad \Delta = p\,. \tag{5.39}$$

For these multiplets the dimensions are

$$\begin{aligned} \dim \mathcal{B}^{\frac{1}{2},\frac{1}{2}}_{[0,p,0](0,0)} &= 2^8\, d(0,p-2,0) = \tfrac{64}{3}\,\hat{p}(\hat{p}-1)^2(\hat{p}-2)\,, \\ \dim \mathcal{B}^{\frac{1}{4},\frac{1}{4}}_{[q,p,q](0,0)} &= 2^{12} d(q-2,p,q) + \tfrac{256}{3}\,\hat{p}(2\hat{p}+1)(2\hat{p}-1)(2\hat{q}+\hat{p}-2)\,. \end{aligned} \tag{5.40}$$

The $\frac{1}{2}$-BPS multiplet corresponds to the chiral primary operators in $\mathcal{N}=4$ theories. The details of the representations for all conformal primary states in the multiplet are well known, being first given in [15] and were described diagrammatically in a fashion similar to that for the $\mathcal{N}=2$ case in [2]. The representations of conformal primary states are obtained, as described above for a long multiplet, from (5.9) or (5.4) and their conjugates. Beyond the simple $\frac{1}{2}$-BPS case the numbers of representations proliferate hugely. We list the self conjugate representations, of the form $[q,p,q]_\ell \equiv [q,p,q]_{(\frac{1}{2}\ell,\frac{1}{2}\ell)}$ which are generated by equal numbers of Q and $\bar{Q}$ supercharges, refered to here as diagonal. This are just the representations that should contribute to the operator product expansion of two chiral primary operators. For the $\frac{1}{2}$-BPS and $\frac{1}{4}$-BPS cases the results for the scale dimension and associated representations are

p	$p+1$	$p+2$	$p+3$	$p+4$
$[0,p,0]_0$	$[1,p-2,1]_1$	$\begin{matrix}[0,p-2,0]_2\\ [2,p-4,2]_0\end{matrix}$	$[1,p-4,1]_1$	$[0,p-4,0]_0$

Table 1. Diagonal representations in $\mathcal{B}^{\frac{1}{2},\frac{1}{2}}_{[0,p,0](0,0)}$.

$2q+p$	$2q+p+1$	$2q+p+2$
$[q,p,q]_0$	$[q{-}1,p,q{-}1]_1$,$[q{-}1,p{+}2,q{-}1]_1$ $2[q,p,q]_1$,$[q{+}1,p{-}2,q{+}1]_1$	$[q{-}2,p{+}2,q{-}2]_2$,$2[q{-}1,p,q{-}1]_2$,$[q,p{-}2,q]_2$,$[q,p,q]_2$ $[q{-}2,p,q{-}2]_0$,$[q{-}2,p{+}2,q{-}2]_0$,$[q{-}2,p{+}4,q{-}2]_0$ $2[q{-}1,p,q{-}1]_0$,$2[q{-}1,p{+}2,q{-}1]_0$,$[q,p{-}2,q]_0$,$3[q,p,q]_0$ $2[q{+}1,p{-}2,q{+}1]_0$,$[q{+}2,p{-}4,q{+}2]_0$

$2q+p+3$	$2q+p+4$
$[q{-}1,p,q{-}1]_3$,$[q{-}3,p{+}2,q{-}3]_1$,$[q{-}3,p{+}4,q{-}3]_1$ $2[q{-}2,p,q{-}2]_1$,$4[q{-}2,p{+}2,q{-}2]_1$,$[q{-}1,p{-}2,q{-}1]_1$ $6[q{-}1,p,q{-}1]_1$,$[q{-}1,p{+}2,q{-}1]_1$,$4[q,p{-}2,q]_1$ $2[q,p,q]_1$,$[q{+}1,p{-}4,q{+}1]_1$,$[q{+}1,p{-}2,q{+}1]_1$	$[q{-}2,p,q{-}2]_2$,$[q{-}2,p{+}2,q{-}2]_2$,$2[q{-}1,p,q{-}1]_2$ $[q,p{-}2,q]_2$,$[q{-}4,p{+}4,q{-}4]_0$,$2[q{-}3,p{+}2,q{-}3]_0$ $3[q{-}2,p,q{-}2]_0$,$[q{-}2,p{+}2,q{-}2]_0$,$2[q{-}1,p{-}2,q{-}1]_0$ $2[q{-}1,p,q{-}1]_0$,$[q,p{-}4,q]_0$,$[q,p{-}2,q]_0$,$[q,p,q]_0$

$2q+p+5$	$2q+p+6$
$[q{-}3,p{+}2,q{-}3]_1$,$2[q{-}2,p,q{-}2]_1$,$[q{-}1,p{-}2,q{-}1]_1$,$[q{-}1,p,q{-}1]_1$	$[q-2,p,q-2]_0$

Table 2. Diagonal representations in $\mathcal{B}^{\frac{1}{4},\frac{1}{4}}_{[q,p,q](0,0)}$.

If $p < 4$ or $p, q < 4$ there are cancellations which will be considered later.

If both the semi-short conditions (5.17) and (5.30) are applied we obtain a semi-short supermultiplet $\mathcal{C}^{t,\bar{t}}_{[k,p,q](j,\bar{j})}$ for various $t, \bar{t}$. The main possible examples, with scale dimensions for the lowest weight states, are given by

$$\begin{aligned}
&\mathcal{C}^{\frac{1}{4},\frac{1}{4}}_{[k,p,q](j,\bar{j})}\,, \qquad k-q = 2(\bar{j}-j)\,, \qquad \Delta = 2+j+\bar{j}+k+p+q\,,\\
&\mathcal{C}^{\frac{1}{2},\frac{1}{2}}_{[0,p,0](j,j)}\,, \qquad\qquad\qquad\quad \Delta = 2+2j+p\,,\\
&\mathcal{C}^{\frac{1}{4},\frac{1}{2}}_{[k,p,0](j,\bar{j})}\,, \qquad k = 2(\bar{j}-j)\,, \qquad \Delta = 2+j+\bar{j}+k+p\,,\\
&\mathcal{C}^{\frac{1}{4},\frac{3}{4}}_{[k,0,0](j,\bar{j})}\,, \qquad k = 2(\bar{j}-j)\,, \qquad \Delta = 2+j+\bar{j}+k\,,\\
&\mathcal{C}^{1,1}_{[0,0,0](j,j)}\,, \qquad\qquad\qquad\quad \Delta = 2+2j\,,
\end{aligned} \tag{5.41}$$

where other cases are obtained under conjugation $\mathcal{C}^{t,\bar{t}}_{[k,p,q](j,\bar{j})} \leftrightarrow \mathcal{C}^{\bar{t},t}_{[q,p,k](\bar{j},j)}$. For the dimension of $\mathcal{C}^{\frac{1}{4},\frac{1}{4}}$ we have

$$\begin{aligned}
\dim \mathcal{C}^{\frac{1}{4},\frac{1}{4}}_{[k,p,q](j,\bar{j})} &= a(k,p,q)\mathcal{J}\bar{\mathcal{J}} + b(k,p,q)\mathcal{J} + b(q,p,k)\bar{\mathcal{J}} + c(k,p,q)\,,\\
a(k,p,q) &= 2^{14} d(k-\tfrac{1}{2}, p, q-\tfrac{1}{2})\\
&\quad - \tfrac{256}{3}\,\hat{p}(\hat{k}+\hat{p}+\hat{q}-1)\big(4\hat{k}^2+4\hat{q}^2+8\hat{k}\hat{p}+8\hat{p}\hat{q}+8\hat{k}\hat{q}-8\hat{k}-8\hat{p}-8\hat{q}+1\big)\,,\\
b(k,p,q) &= 2^{13} d(k-\tfrac{1}{2}, p, q)\\
&\quad - \tfrac{256}{3}\,\hat{p}\big(4\hat{k}^3+2\hat{q}^3+12\hat{k}^2\hat{p}+8\hat{k}\hat{p}^2+12\hat{k}^2\hat{q}+6\hat{k}\hat{q}^2+4\hat{p}^2\hat{q}+6\hat{p}\hat{q}^2+18\hat{k}\hat{p}\hat{q}\\
&\qquad\qquad - 6\hat{k}^2-4\hat{p}^2-3\hat{q}^2-12\hat{k}\hat{p}-12\hat{k}\hat{q}-9\hat{p}\hat{q}+1\big)\,,\\
c(k,p,q) &= 2^{12} d(k,p,q) - \tfrac{256}{3}\,\hat{p}(\hat{k}+\hat{p}+\hat{q})\big(2\hat{k}^2+2\hat{q}^2+4\hat{k}\hat{p}+4\hat{p}\hat{q}+4\hat{k}\hat{q}-3\big)\,.
\end{aligned} \tag{5.42}$$

The dimensions for other examples may be obtained as special cases of (5.42), thus $\dim \mathcal{C}^{\frac{1}{4},\frac{1}{2}}_{[k,p,0](j,\bar{j})}$ is obtained by setting $q = 0$. In particular we may obtain

$$\dim \mathcal{C}^{\frac{1}{2},\frac{1}{2}}_{[0,p,0](j,j)} = 2^{12} d(0,p-1,0)\mathcal{J}(\mathcal{J}+2) + \tfrac{256}{3}\,\hat{p}\,(2\hat{p}-1)(2\hat{p}+1)(\hat{p}+2\mathcal{J}+2)\,. \tag{5.43}$$

For $\mathcal{C}^{1,1}_{[0,0,0](j,\bar{j})}$, relaxing the constraint $j = \bar{j}$, we have from imposing both (5.17) and (5.30), with the basis defined by (5.1),

$$\big(4j\bar{j}\,\mathrm{P}_{21} + 2j\,\mathrm{P}_{22}\,J_- - 2\bar{j}\mathrm{P}_{11}\,\bar{J}_- - \mathrm{P}_{12}\,J_-\bar{J}_-\big)|0,0,0;j,\bar{j}\rangle^{\mathrm{hw}} = 0\,, \tag{5.44}$$

which is equivalent to the conservation equation (4.41). Using (5.24) we may easily con-

struct the full supermultiplet in this case,



(5.45)

For $j, \bar{\jmath} > 0$ all representations correspond to conserved currents. It is easy to see that

$$\dim \mathcal{C}^{1,1}_{[0,0,0](j,\bar{\jmath})} = 2^8(\mathcal{J} + \bar{\mathcal{J}} + 3)\,. \tag{5.46}$$

For $j = \bar{\jmath} = 0$ this would correspond to the Konishi scalar in free theory, all states obtained for $l, \bar{l} = 1, 2, 3, 4$ satisfy conservation equations, for $l = \bar{l} = 1$ there is a conserved vector curent. Such higher spin conserved currents (for $j = \bar{\jmath} = \frac{1}{2}\ell$ they are described by symmetric traceless rank ℓ tensors) are found in free field theories but should be absent in interacting theories[1]. There is a crucial difference with $\mathcal{N} = 2$ for $j = \bar{\jmath} = 0$, described

[1] Higher spin currents in conformal theories have been considered by Anselmi [24] and also in [25].

by (4.40), since the maximal spin in that case corresponds to the the conserved energy momentum tensor and there are no higher spin currents.

As earlier we list the self-conjugate representations obtained by the action of equal numbers of Q and $\bar{Q}$ supercharges

$\ell+p+2$	$\ell+p+3$	$\ell+p+4$
$[0,p,0]_\ell$	$2[0,p,0]_{\ell+1}$,$[1,p-2,1]_{\ell+1}$ $[1,p,1]_{\ell+1}$,$[1,p-2,1]_{\ell-1}$	$[0,p-2,0]_{\ell+2}$,$3[0,p,0]_{\ell+2}$,$[0,p+2,0]_{\ell+2}$,$2[1,p-2,1]_{\ell+2}$ $2[1,p,1]_{\ell+2}$,$[2,p-2,2]_{\ell+2}$,$[0,p-2,0]_\ell$ $[0,p,0]_\ell$,$4[1,p-2,1]_\ell$,$[2,p-4,2]_\ell$,$[2,p-2,2]_\ell$ $[0,p-2,0]_{\ell-2}$

$\ell+p+5$	$\ell+p+6$
$2[0,p,0]_{\ell+3}$,$[1,p-2,1]_{\ell+3}$,$[1,p,1]_{\ell+3}$ $2[0,p-2,0]_{\ell+1}$,$2[0,p,0]_{\ell+1}$,$[1,p-4,1]_{\ell+1}$,$6[1,p-2,1]_{\ell+1}$ $[1,p,1]_{\ell+1}$,$2[2,p-4,2]_{\ell+1}$,$2[2,p-2,2]_{\ell+1}$,$[3,p-4,3]_{\ell+1}$ $2[0,p-2,0]_{\ell-1}$,$[1,p-4,1]_{\ell-1}$,$[1,p-2,1]_{\ell-1}$	$[0,p,0]_{\ell+4}$,$[0,p-2,0]_{\ell+2}$,$[0,p,0]_{\ell+2}$ $4[1,p-2,1]_{\ell+2}$,$[2,p-4,2]_{\ell+2}$,$[2,p-2,2]_{\ell+2}$ $[0,p-4,0]_\ell$,$3[0,p-2,0]_\ell$,$[0,p,0]_\ell$ $2[1,p-4,1]_\ell$,$2[1,p-2,1]_\ell$,$[2,p-4,2]_\ell$

$\ell+p+7$	$\ell+p+8$
$[1,p-2,1]_{\ell+3}$,$2[0,p-2,0]_{\ell+1}$ $[1,p-4,1]_{\ell+1}$,$[1,p-2,1]_{\ell+1}$	$[0,p-2,0]_{\ell+2}$

Table 3. Diagonal representations in $\mathcal{C}^{\frac{1}{2},\frac{1}{2}}_{[0,p,0](\frac{1}{2}\ell,\frac{1}{2}\ell)}$.

The rather more lengthy result for $\mathcal{C}^{\frac{1}{4},\frac{1}{4}}_{[q,p,q](\frac{1}{2}\ell,\frac{1}{2}\ell)}$ is given in appendix C.

If we apply the semi-shortening conditions (5.6) together with (5.29), so that $j=0$, we obtain supermultiplets denoted here by $\mathcal{D}^{s,\bar{t}}_{[k,p,q](0,\bar{\jmath})}$. The main case is

$$\mathcal{D}^{\frac{1}{4},\frac{1}{4}}_{[k,p,q](0,\bar{\jmath})}\,, \qquad k-q=2+2\bar{\jmath}\,, \qquad \Delta=1+\bar{\jmath}+k+p+q\,. \tag{5.47}$$

We may also allow $\bar{t}=\frac{1}{2},\frac{3}{4}$ if $q=0, p=q=0$ respectively but $s=\frac{1}{2},\frac{3}{4}$ are not possible. Of course there are also the conjugate supermultiplets $\bar{\mathcal{D}}^{t,\bar{s}}_{[k,p,q](j,0)}$ for $\bar{s}=\frac{1}{4}$, $t=\frac{1}{4},\frac{1}{2},\frac{3}{4}$. We have

$$\begin{aligned}\dim \mathcal{D}^{\frac{1}{4},\frac{1}{4}}_{[k,p,q](0,\bar{\jmath})} &= b(q,p,k-1)\bar{\mathcal{J}}+c(k-1,p,q)\,,\\ \dim \bar{\mathcal{D}}^{\frac{1}{4},\frac{1}{4}}_{[k,p,q](j,0)} &= b(k,p,q-1)\mathcal{J}+c(k,p,q-1)\,,\end{aligned} \tag{5.48}$$

with b,c given in (5.42), and

$$\begin{aligned}\dim \bar{\mathcal{D}}^{\frac{1}{2},\frac{1}{4}}_{[0,p,q](j,0)} &= 2^{12}d(0,p,q-1)(\mathcal{J}+1)\\ &\quad -\tfrac{256}{3}\hat{p}\big((2\hat{p}+2\hat{q}-1)(4\hat{p}\hat{q}+2\hat{q}^2-2\hat{p}-2\hat{q}-1)\mathcal{J}+(\hat{p}+\hat{q})(4\hat{p}\hat{q}+2\hat{q}^2-3)\big)\,,\\ \dim \bar{\mathcal{D}}^{\frac{3}{4},\frac{1}{4}}_{[0,0,q](j,0)} &= 2^{11}d(0,0,q-1)(\mathcal{J}+\tfrac{3}{2})-256\big((2\hat{q}^2-1)\mathcal{J}+\hat{q}\big)\,.\end{aligned} \tag{5.49}$$

In (5.42) or (5.48) and (5.49) we have not imposed the relations which determine $\mathcal{J}-\bar{\mathcal{J}}$ or $\mathcal{J}$ in terms of k,p,q given in (5.41) or (5.47).

Just as for $\mathcal{N}=2$ we may decompose the long supermultiplet $\mathcal{A}^{\Delta}_{[k,p,q](j,\bar{j})}$ at the unitarity threshold given by (5.37). When both conditions in (5.37) hold simultaneously we have, extending (5.34), the semi-direct sum

$$\begin{aligned}
&\mathcal{A}^{2-j+3\bar{j}+p+2q}_{[k,p,q](j,\bar{j})}\big|_{k-q=2(\bar{j}-j)} \\
&\qquad\simeq \mathcal{C}^{\frac14,\frac14}_{[k,p,q](j,\bar{j})}\oplus\mathcal{C}^{\frac14,\frac14}_{[k+1,p,q](j-\frac12,\bar{j})}\oplus\mathcal{C}^{\frac14,\frac14}_{[k,p,q+1](j,\bar{j}-\frac12)}\oplus\mathcal{C}^{\frac14,\frac14}_{[k+1,p,q+1](j-\frac12,\bar{j}-\frac12)}\,,\\
&\mathcal{A}^{2+2j+p}_{[0,p,0](j,j)}\simeq \mathcal{C}^{\frac12,\frac12}_{[0,p,0](j,j)}\oplus\mathcal{C}^{\frac14,\frac12}_{[1,p,0](j-\frac12,j)}\oplus\mathcal{C}^{\frac12,\frac14}_{[0,p,1](j,j-\frac12)}\oplus\mathcal{C}^{\frac14,\frac14}_{[1,p,1](j-\frac12,j-\frac12)}\,,\\
&\mathcal{A}^{2+2j}_{[0,0,0](j,j)}\simeq \mathcal{C}^{1,1}_{[0,0,0](j,j)}\oplus\mathcal{C}^{\frac14,\frac34}_{[1,0,0](j-\frac12,j)}\oplus\mathcal{C}^{\frac34,\frac14}_{[0,0,1](j,j-\frac12)}\oplus\mathcal{C}^{\frac14,\frac14}_{[1,0,1](j-\frac12,j-\frac12)}\,,
\end{aligned}\tag{5.50}$$

as well as, for $j=\bar{j}=0$,

$$\begin{aligned}
&\mathcal{A}^{2+p+2q}_{[q,p,q](0,0)}\simeq \mathcal{C}^{\frac14,\frac14}_{[q,p,q](0,0)}\oplus\mathcal{D}^{\frac14,\frac14}_{[q+2,p,q](0,0)}\oplus\bar{\mathcal{D}}^{\frac14,\frac14}_{[q,p,q+2](0,0)}\oplus\mathcal{B}^{\frac14,\frac14}_{[q+2,p,q+2](0,0)}\,,\\
&\mathcal{A}^{2+p}_{[0,p,0](0,0)}\simeq \mathcal{C}^{\frac12,\frac12}_{[0,p,0](0,0)}\oplus\mathcal{D}^{\frac14,\frac12}_{[2,p,0](0,0)}\oplus\bar{\mathcal{D}}^{\frac12,\frac14}_{[0,p,2](0,0)}\oplus\mathcal{B}^{\frac14,\frac14}_{[2,p,2](0,0)}\,,\\
&\mathcal{A}^{2}_{[0,0,0](0,0)}\simeq \mathcal{C}^{1,1}_{[0,0,0](0,0)}\oplus\mathcal{D}^{\frac14,\frac34}_{[2,0,0](0,0)}\oplus\bar{\mathcal{D}}^{\frac34,\frac14}_{[0,0,2](0,0)}\oplus\mathcal{B}^{\frac14,\frac14}_{[2,0,2](0,0)}\,.
\end{aligned}\tag{5.51}$$

Note that, corresponding to (5.35), we may identify

$$\mathcal{C}^{\frac14,\frac14}_{[k,p,q](j,-\frac12)}\simeq\bar{\mathcal{D}}^{\frac14,\frac14}_{[k,p,q+1](j,0)}\,,\qquad \mathcal{C}^{\frac14,\frac14}_{[k,p,q](-\frac12,-\frac12)}\simeq\mathcal{B}^{\frac14,\frac14}_{[k+1,p,q+1](0,0)}\,,\tag{5.52}$$

which ensures that (5.51) is a special case of (5.50).

For $\mathcal{N}=4$ it is also of interest to consider the central extension which is obtained by letting in (3.4a,b) $R^i{}_j\to R^i{}_j+\frac12\delta^i{}_jZ$ when the superconformal group is $SU(2,2|4)$. The multiplet structure is unchanged but in (5.8), (5.17) $\Delta\to\Delta-Z$ whereas in (5.28), (5.30) $\Delta\to\Delta+Z$. For short supermultiplets $\mathcal{B}^{s,\bar{s}}_{[k,p,q](0,0)}$ we no longer require $s=\bar{s}$ or $k=q$ if $s=\bar{s}=\frac14$. As well as (5.38) and (5.39) we may consider

$$\mathcal{B}^{\frac14,\frac34}_{[k,0,0](0,0)}\,,\quad \Delta=k\,,\quad Z=-\tfrac12k\,,\qquad \mathcal{B}^{\frac34,\frac14}_{[0,0,q](0,0)}\,,\quad \Delta=q\,,\quad Z=\tfrac12q\,,\tag{5.53}$$

with

$$\dim\mathcal{B}^{\frac14,\frac34}_{[k,0,0](0,0)}=2^8d(k-2,0,0)\,,\qquad \dim\mathcal{B}^{\frac34,\frac14}_{[0,0,q](0,0)}=2^8d(0,0,q-2)\,.\tag{5.54}$$

For the semi-short multiplets $\mathcal{C}^{t,\bar{t}}_{[k,p,q](j,\bar{j})}$ in (5.41) there is no longer any constraint on $\bar{j}-j$ although the formulae for Δ are unchanged. The semi-short/short multiplet $\mathcal{D}^{s,\bar{t}}_{[k,p,q](0,\bar{j})}$,

and its conjugate $\mathcal{D}^{\bar{s},t}_{[k,p,q](j,0)}$, are also no longer restricted to $s, \bar{s} = \frac{1}{4}$. For $\mathcal{D}^{\frac{1}{4},\frac{1}{4}}_{[k,p,q](0,\bar{j})}$ there is now no constraint on $k-q$ as in (5.47) although Δ is the same. Of particular interest are the chiral multiplets $\mathcal{D}^{1,1}_{[0,0,0](0,\bar{j})}$ and $\bar{\mathcal{D}}^{1,1}_{[0,0,0](j,0)}$. For the latter case consistency of (5.15) and (5.26) for $i = 1, 2, 3, 4$ with (3.1) leads to the Dirac equation, similarly to (4.45), which takes the form

$$\big(2j\,\mathrm{P}_{2\dot{\alpha}} - \mathrm{P}_{1\dot{\alpha}}\,J_-\big)|0,0,0;j,0\rangle^{\mathrm{hw}} = 0\,, \tag{5.55}$$

where the state $|0,0,0;j,0\rangle^{\mathrm{hw}}$ has dimension $\Delta = 1 + j$.

6. Applications in Special Cases

In the previous two sections we gave results for constructing supermultiplets by the action of supercharges on the lowest dimension state when this was assumed to belong to a general representation of $SU(2)_R$ or $SU(4)_R$. In such a generic case the Dynkin labels for the resulting representations are obtained simply by adding the weights of the relevant supercharges to Dynkin indices for the representation to which the lowest dimension state belongs. For particular examples the indices obtained are not necessarily positive or zero. In such circumstances the Racah Speiser algorithm, described more fully in appendix B, provides a straightforward procedure for dealing with these situations.

We first consider the $\mathcal{N} = 2$ case. Identifying representations with their labels $R_{(j,\bar{j})}$ as in section 4 the algorithm requires that in tensor product expansions we should identify

$$(-R)_{(j,\bar{j})} = -(R-1)_{(j,\bar{j})}\,, \qquad R_{(-j,\bar{j})} = -R_{(j-1,\bar{j})}\,, \qquad R_{(j,-\bar{j})} = -R_{(j,\bar{j}-1)}\,. \tag{6.1}$$

Necessarily in a representation $R_{(j,\bar{j})}$ if any of $R, j, \bar{j}$ is equal to $-\frac{1}{2}$ the contribution of this representation in a tensor product expansion must be set to zero. Results following from (6.1) have been used implicitly at various points in section 4. As an illustration we may consider (4.15) for $R = \frac{1}{2}$ when there is a further truncation,

$$\tfrac{1}{2}{}_{(0,\bar{j})} \xrightarrow{Q} 0_{(\frac{1}{2},\bar{j})}\,, \qquad \tfrac{1}{2}{}_{(j,0)} \xrightarrow{\bar{Q}} 0_{(j,\frac{1}{2})}\,. \tag{6.2}$$

If we apply the results to the short multiplet $\hat{\mathcal{B}}_R$, illustrated in (4.22), for $R = \frac{1}{2}, 1$ we find



(6.3)

The negative contributions may be cancelled by imposing the equations of motion or conservation equations. Thus in the $R = \frac{1}{2}$ case representing the contribution with $\Delta = 1$ by a scalar field φ^i and for $\Delta = \frac{3}{2}$ by two fermions ψ_α, $\chi^{\dot{\alpha}}$ it is easy to see that $\partial^2\varphi^i$ and $\partial^{\dot{\alpha}\alpha}\psi_\alpha$, $\partial_{\alpha\dot{\alpha}}\chi^{\dot{\alpha}}$ exactly correspond in dimensions and spins to the negative representations with $\Delta = 3$ and $\Delta = \frac{5}{2}$ respectively. Thus the requirement of the equations of motion cancels these negative terms and this represents the usual $\mathcal{N} = 2$ hypermultiplet. In the case of $\hat{\mathcal{B}}_1$ the multiplet contains a $SU(2)_R$ singlet current $J_{\alpha\dot{\alpha}}$ with $\Delta = 3$, belonging to the representation $0_{(\frac{1}{2},\frac{1}{2})}$, for which the divergence $\partial^{\dot{\alpha}\alpha}J_{\alpha\dot{\alpha}}$ corresponds to the negative contribution in (6.3) and so the conservation of the current cancels the negative piece. This case is the simplest gauge invariant short multiplet in $\mathcal{N} = 2$ theories.

For the semi-short modules we may note, using (6.1), that (4.27) reduces to (4.28) for $R = 0$. The semi-short supermultiplet $\mathcal{C}_{R(j,\bar{j})}$, displayed in (4.37) reduces to $\mathcal{C}_{0(j,\bar{j})}$, as given in (4.40), together with negative contributions which are cancelled by imposing conservation equations on all fields belonging to the supermultiplet. We may also note that in (4.27) setting $j = -\frac{1}{2}$ removes the first term and this module is identical to the short module exhibited in (4.15) for $R \to R + \frac{1}{2}$. This provides an explanation of (4.51).

A further special case, following from (6.2), is obtained by considering (4.43) for $R = \frac{1}{2}$,



(6.4)

neglecting negative terms which are removed by imposing conservation of the currents for the representations $0_{(\frac{1}{2},\bar{\jmath})}$, if $\bar{\jmath} > 0$, $\frac{1}{2}{}_{(\frac{1}{2},\bar{\jmath}+\frac{1}{2})}$ and $0_{(\frac{1}{2},\bar{\jmath}+1)}$. In addition we may obtain from (4.37) and (4.22), using (6.1),

$$\mathcal{C}_{R(-1,-1)} = \hat{\mathcal{B}}_{R(0,0)}\,. \tag{6.5}$$

A similar pattern emerges for $\mathcal{N} = 4$. In this case for $SU(4)$ representations $[k,p,q]$ in tensor product expansions we should identify

$$\begin{aligned} [k,p,q] = &-[-k-2,p+k+1,q]\,, \qquad [k,p,q] = -[k+p+1,-p-2,q+p+1]\,, \\ &[k,p,q] = -[k,p+q+1,-q-2]\,. \end{aligned} \tag{6.6}$$

These identities generate a group of order 24 so there are 23 non trivial relations which allow $[k,p,q] = \pm[k',p',q']$ with $k',p',q' \geq 0$ or $[k,p,q] = 0$, as when any of k,p,q are equal to -1. These rules allow the construction of supermultiplets described previously to be extended to cases when the lowest state representation has low k,p,q. Some examples are given in tables in appendix C.

Using (6.6) it is easy to see that there is also a truncation of the basic short modules for special choices of k,p,q. For $b^{\frac{1}{4}}_{[k,p,q]}$, given by (5.9), if $k = 1$

$$\begin{aligned} [1,p,q]_{(0,\bar{\jmath})} &\xrightarrow{Q} \begin{matrix}[1,p,q-1],[1,p-1,q+1]_{(\frac{1}{2},\bar{\jmath})} \\ [0,p+1,q]_{(\frac{1}{2},\bar{\jmath})}\end{matrix} \xrightarrow{Q^2} \begin{matrix}[1,p-1,q],[0,p+1,q-1],[0,p,q+1]_{(1,\bar{\jmath})} \\ [1,p,q-2],[1,p-1,q],[1,p-2,q+2],[0,p+1,q-1],[0,p,q+1]_{(0,\bar{\jmath})}\end{matrix} \\ &\xrightarrow{Q^3} \begin{matrix}[0,p,q]_{(\frac{3}{2},\bar{\jmath})},[1,p-1,q-1],[1,p-2,q+1]_{(\frac{1}{2},\bar{\jmath})} \\ [0,p+1,q-2],2[0,p,q],[0,p-1,q+2]_{(\frac{1}{2},\bar{\jmath})}\end{matrix} \xrightarrow{Q^4} \begin{matrix}[0,p,q-1],[0,p-1,q+1]_{(1,\bar{\jmath})} \\ [1,p-2,q],[0,p,q-1],[0,p-1,q+1]_{(0,\bar{\jmath})}\end{matrix} \\ &\xrightarrow{Q^5} [0,p-1,q]_{(\frac{1}{2},\bar{\jmath})}\,. \end{aligned} \tag{6.7}$$

For $b^{\frac{1}{2}}_{[0,p,q]}$, given by (5.11), we have if $p=1$

$$[0,1,q]_{(0,\bar{j})} \xrightarrow{Q} \begin{matrix}[0,1,q-1]_{(\frac{1}{2},\bar{j})}\\ [0,0,q+1]_{(\frac{1}{2},\bar{j})}\end{matrix} \xrightarrow{Q^2} \begin{matrix}[0,0,q]_{(1,\bar{j})}\\ [0,1,q-2],[0,0,q]_{(0,\bar{j})}\end{matrix} \xrightarrow{Q^3} [0,0,q-1]_{(\frac{1}{2},\bar{j})}\,, \tag{6.8}$$

and for $b^{\frac{3}{4}}_{[0,0,q]}$, given by (5.13), for $q=1$ we have just,

$$[0,0,1]_{(0,\bar{j})} \xrightarrow{Q} [0,0,0]_{(\frac{1}{2},\bar{j})}\,. \tag{6.9}$$

In each example there is one less level than for the general case. In (6.7) there is one further stage of shortening if $p=0$ and in (6.8) if $q=0$. In general for these modules we have

$$b^{\frac{1}{4}}_{[0,p,q]} \simeq b^{\frac{1}{2}}_{[0,p,q]} \ominus b^{\frac{1}{2}}_{[0,p+1,q]}\,, \qquad b^{\frac{1}{2}}_{[0,0,q]} \simeq b^{\frac{3}{4}}_{[0,0,q]}\,. \tag{6.10}$$

For the semi-short modules $c^{t}_{[k,p,q]j}$ there is no similar truncation but it easy to check, with the aid of results flowing from (6.6), that these are nested in that $c^{\frac{1}{4}}_{[0,p,q]j} \simeq c^{\frac{1}{2}}_{[0,p,q]j}$ and similarly if p and then q are set to zero for $t=\frac{3}{4},1$. Thus (5.19) and (5.21) are identical for $k=0$. We may also note that in (5.18) if $j=-\frac{1}{2}$ the first representation is absent and the module is identical with (5.9) with $k\to k+1$, justifying (5.35). In a similar fashion we may note that

$$c^{\frac{1}{2}}_{[0,p,q]-1} \simeq \ominus b^{\frac{1}{2}}_{[0,p,q]} \oplus b^{\frac{1}{2}}_{[0,p+1,q]}\,. \tag{6.11}$$

These results extend to complete supermultiplets. The simplest non trivial $\frac{1}{2}$-BPS multiplets are $\mathcal{B}^{\frac{1}{2},\frac{1}{2}}_{[0,1,0](0,0)}$, which has maximal shortening,



(6.12)

and $\mathcal{B}^{\frac{1}{2},\frac{1}{2}}_{[0,2,0](0,0)}$



(6.13)

In (6.12), as in the $\mathcal{N} = 2$ case, the negative contributions are cancelled by imposing equations of motion on all fields (for the representation $[0, 0, 0]_{(1,0)}$ with $\Delta = 2$ corresponding to a field $f_{\alpha\beta} = f_{\beta\alpha}$ there is associated vector $V_{\alpha\dot{\alpha}}$, $[0, 0, 0]_{(\frac{1}{2},\frac{1}{2})}$, with $\Delta = 3$ which contributes negatively and a scalar φ, $[0, 0, 0]_{(0,0)}$, with $\Delta = 4$, the equation of motion is $\partial^{\dot{\alpha}\alpha} f_{\alpha\beta} = 0$ and we may note that $f \xrightarrow{\partial} V \xrightarrow{\partial} \varphi$ with $\partial\partial f = 0$). For each column there are zero total degrees of freedom reflecting the vanishing of the dimension formula (5.40) in this case. This of course corresponds to the elementary multiplet in $\mathcal{N} = 4$ gauge theories. In (6.13) it is necessary to impose conservation equations for the $[1, 0, 1]_{(\frac{1}{2},\frac{1}{2})}$ $SU(4)$ current, as well as the energy momentum tensor, $[0, 0, 0]_{(1,1)}$, and spinor currents, $[1, 0, 0]_{(1,\frac{1}{2})}$ and $[0, 0, 1]_{(\frac{1}{2},1)}$. For $p > 2$ there are no such constraints on the corresponding fields since no negative representations appear in $\mathcal{B}^{\frac{1}{2},\frac{1}{2}}_{[0,p,0](0,0)}$. As a consequence of (6.7) we may also note that $\mathcal{B}^{\frac{1}{4},\frac{1}{4}}_{[1,p,1](0,0)}$ is shortened from the generic $\frac{1}{4}$-BPS case, the scale dimension ranges just from $p + 2$ to $p + 7$. Using (6.7) we also obtain the singleton multiplet $\mathcal{B}^{\frac{1}{4},\frac{3}{4}}_{[1,0,0](0,0)}$ is

described by



(6.14)

with all fields required to satisfy equations of motion. The singleton representations $\mathcal{B}^{\frac{1}{2},\frac{1}{2}}_{[0,1,0](0,0)}$, $\mathcal{B}^{\frac{1}{4},\frac{3}{4}}_{[1,0,0](0,0)}$ and $\mathcal{B}^{\frac{3}{4},\frac{1}{4}}_{[0,0,1](0,0)}$ are building blocks for all supermultiplets. We may also note that $\mathcal{B}^{\frac{1}{4},\frac{3}{4}}_{[2,0,0](0,0)}$ contains conserved currents for the representations $[0,1,0]_{(\frac{1}{2},\frac{1}{2})}$, $[0,0,1]_{(1,\frac{1}{2})}$ and $[0,0,0]_{(\frac{3}{2},\frac{1}{2})}$.

For the semi-short multiplets $\mathcal{C}^{t,\bar{t}}_{[k,p,q](j,\bar{j})}$ we have verified that $\mathcal{C}^{\frac{1}{4},\frac{1}{4}}_{[0,p,0](j,\bar{j})} \simeq \mathcal{C}^{\frac{1}{2},\frac{1}{2}}_{[0,p,0](j,\bar{j})}$ and also $\mathcal{C}^{\frac{1}{2},\frac{1}{2}}_{[0,0,0](j,\bar{j})}$ reduces to $\mathcal{C}^{1,1}_{[0,0,0](j,\bar{j})}$, shown in (5.45), together with negative terms corresponding to the divergences of all currents with spins $j, \bar{j} > 0$. In a similar fashion we may show that $\bar{\mathcal{D}}^{\frac{1}{2},\frac{1}{2}}_{[0,1,0](j,0)}$ contains higher spin conserved currents with the representations $[1,0,0]_{(j+\frac{1}{2},\frac{1}{2}),(j+\frac{1}{2},1),(j+2,\frac{1}{2})}$, $[0,1,0]_{(j+\frac{1}{2},\frac{1}{2}),(j+1,\frac{1}{2})}$, $[2,0,0]_{(j+\frac{1}{2},\frac{1}{2})}$, $[0,0,0]_{(j,1),(j+\frac{3}{2},\frac{1}{2}),(j+2,1)}$, $[0,0,1]_{(j+1,\frac{1}{2}),(j+\frac{3}{2},1)}$, $[1,0,1]_{(j+\frac{3}{2},\frac{1}{2})}$ and $[1,1,0]_{(j+1,\frac{1}{2})}$, in each case with scale dimension $j+\bar{j}+2$ as is necessary for consistency with conformal invariance. The relation (5.52) demonstrates that semi-short multiplets are connected to short multiplets in special cases, in addition from (6.11) we may note that

$$\mathcal{C}^{\frac{1}{2},\frac{1}{2}}_{[0,p,0](-1,-1)} \simeq \mathcal{B}^{\frac{1}{2},\frac{1}{2}}_{[0,p,0](0,0)} \ominus 2\mathcal{B}^{\frac{1}{2},\frac{1}{2}}_{[0,p+1,0](0,0)} \oplus \mathcal{B}^{\frac{1}{2},\frac{1}{2}}_{[0,p+2,0](0,0)}\,. \tag{6.15}$$

Allowing for a central extension there are several cases leading to quite small multiplets. These have been discussed in [16][2].

[2] In detail for the first paper in [16] Table 1 corresponds to $\mathcal{B}^{\frac{1}{2},\frac{1}{2}}_{[0,1,0](0,0)}$, Table 2 to $\mathcal{B}^{\frac{1}{4},\frac{3}{4}}_{[1,0,0](0,0)}$, Table 3 to $\mathcal{B}^{\frac{3}{4},\frac{1}{4}}_{[0,0,1](0,0)}$, Table 4 to $\bar{\mathcal{D}}^{1,1}_{[0,0,0](0,0)}$, Table 5 to $\mathcal{D}^{1,1}_{[0,0,0](0,0)}$, Table 6 to $\bar{\mathcal{D}}^{1,1}_{[0,0,0](j-1,0)}$, Table 7 to $\mathcal{D}^{1,1}_{[0,0,0](0,j-1)}$, Table 8 to $\mathcal{B}^{\frac{1}{4},\frac{1}{2}}_{[1,1,0](0,0)}$, Table 9 to $\mathcal{B}^{\frac{1}{2},\frac{1}{4}}_{[0,1,1](0,0)}$, Table 10 to $\bar{\mathcal{D}}^{\frac{1}{2},\frac{1}{2}}_{[0,1,0](j-1,0)}$, Table 11 to $\mathcal{D}^{\frac{1}{2},\frac{1}{2}}_{[0,1,0](0,j-1)}$, Table 12 to $\mathcal{C}^{1,1}_{[0,0,0](j_L-1,j_R-1)}$, while Table 13 corresponds to $\mathcal{B}^{\frac{1}{2},\frac{1}{2}}_{[0,2,0](0,0)}$. In the second paper among the new multiplets Table 6 corresponds to $\mathcal{B}^{\frac{1}{4},\frac{3}{4}}_{[2,0,0](0,0)}$, Table 7 to $\mathcal{B}^{\frac{3}{4},\frac{1}{4}}_{[0,0,2](0,0)}$, Table 8 to $\mathcal{B}^{0,1}_{[0,0,0](0,0)}$ and Table 9 to $\mathcal{B}^{1,0}_{[0,0,0](0,0)}$.

7. Discussion

An initial motivation for this work was to identify supermultiplets with the sets of operators that arise in the operator product expansion analysis of four point functions of the $\mathcal{N} = 4$ superconformal $SU(N)$ gauge theory. In the analysis in [7] of superconformal Ward identities for the four point function for $[0, 2, 0]$ chiral primary operators we showed how the solution of these identities required the presence of sets of operators in the operator product expansion with related scale dimensions and spins. The spectra in each case match exactly with what we have termed diagonal states, formed by equal numbers of Q and $\bar{Q}$ supercharges acting on a superconformal primary state, for particular short and semi-short multiplets[3]. This is as expected according to $U(1)_Y$ bonus symmetry [26,27]. Assuming a non renormalisation theorem such that there is only a single undetermined function of two variables in the solution of the Ward identities then there must be present in the operator product expansion analysis semi-short multiplets $\mathcal{C}^{\frac{1}{2},\frac{1}{2}}_{[0,2,0](j,j)}$, $j = 0, 1, 2\ldots$ and $\mathcal{C}^{\frac{1}{4},\frac{1}{4}}_{[1,0,1](j,j)}$, $j = \frac{1}{2}, \frac{3}{2}, \ldots$. A similar pattern emerges in an extension [28] of this analysis to next extremal [29] four point correlators of chiral primary operators. Assuming non renormalisation theorems the operator product expansion requires the presence of semi-short multiplets $\mathcal{C}^{\frac{1}{2},\frac{1}{2}}_{[0,p,0](j,j)}$ for all $p = 2, 3, \ldots$ and $j = 0, \frac{1}{2}, 1, \ldots$ (a related discussion of next extremal correlators is also given in [6]).

More generally it remains to be understood under what circumstances short or semi-short multiplets are protected against renormalisation effects giving rise to anomalous dimensions. If these are to occur it is necessary the multiplet is part of a possible long multiplet. As a consequence of (5.50) and (5.51) the only multiplets satisfying shortening conditions for the action of both the Q and $\bar{Q}$ supercharges that cannot form part of a long multiplet and therefore must be protected are

$$\mathcal{B}^{\frac{1}{2},\frac{1}{2}}_{[0,p,0](0,0)}\,, \qquad \mathcal{B}^{\frac{1}{4},\frac{1}{4}}_{[1,p,1](0,0)}\,. \tag{7.1}$$

For other cases there is the possibility of joining with other multiplets to form a long multiplet so that such multiplets by themselves cannot be guaranteed to be protected against renormalisation effects. The number of states in the free and interacting theories should be the same but the way in which they are arranged into supermultiplets may change for g non zero. Given the results (5.50) and (5.51) and a knowledge of the free theory supermultiplet struture then, although individual semi-short multiplets can be part of more than one long multiplet, it may be possible, following [22], to set up index theorems

[3] In particular in [7] the states labelled by $\mathcal{B}_\ell$ correspond to $\mathcal{C}^{1,1}_{[0,0,0](\frac{1}{2}\ell-1,\frac{1}{2}\ell-1)}$, $\mathcal{D}_\ell$ to $\mathcal{C}^{\frac{1}{2},\frac{1}{2}}_{[0,2,0](\frac{1}{2}\ell-1,\frac{1}{2}\ell-1)}$, $\mathcal{E}_\ell$ to $\mathcal{C}^{\frac{1}{4},\frac{1}{4}}_{[1,0,1](\frac{1}{2}\ell-1,\frac{1}{2}\ell-1)}$, $\hat{B}_0$ to $\mathcal{B}^{\frac{1}{2},\frac{1}{2}}_{[0,2,0](0,0)}$, $\hat{D}_0$ to $\mathcal{B}^{\frac{1}{2},\frac{1}{2}}_{[0,4,0](0,0)}$ and $\mathcal{E}_0$ to $\mathcal{B}^{\frac{1}{4},\frac{1}{4}}_{[2,0,2](0,0)}$.

which would ensure, for a non zero index, the existence of other protected supermultiplets. On the other hand a $\frac{1}{4}$-BPS multiplet $\mathcal{B}^{\frac{1}{4},\frac{1}{4}}_{[q,p,q](0,0)}$ with $q \geq 2$ can only be part of a unique long multiplet $\mathcal{A}^{p+2q-2}_{[q-2,p,q-2](0,0)}$. At present the knowledge of what protected multiplets may appear in $\mathcal{N}=4$ SYM is rather limited, for a discussion of $\frac{1}{4}$-BPS operators in field theory see [30]. More generally it seems probable that in any interacting theory multiplets containing higher spin currents, such as $\mathcal{C}^{1,1}_{[0,0,0](j,j)}$, for any j, illustrated in (5.45), should be absent (this was implicitly assumed in [7]). This rule of course applies in the simplest case $j=0$ for the Konishi multiplet, the Konishi anomaly affects not just the usual spin 1 vector current but also currents in the same multiplet with spin 2,3,4. In general protected operators belonging to semi-short multiplets are essentially multi-trace operators formed from products of single trace $\frac{1}{2}$-BPS chiral primary operators [31], which are at least protected against perturbative corrections to leading order in N [32].

We may also note that the conditions imposed on the action of the Q and $\bar{Q}$ supercharges, such as (4.24), (4.29) and (4.35) for $\mathcal{N}=2$ and their extensions to $\mathcal{N}=4$, on the superconformal primary states for semi-short multiplets have a direct correspondence in terms of the action of the D and $\bar{D}$ spinor derivatives on superfields [18], where they are referred to as current like protected operators. In the simplest case the $\mathcal{N}=1$ Konishi scalar field $\mathcal{O}$ satisfies $D^2\mathcal{O} = \bar{D}^2\mathcal{O} = 0$ in free theory, analogous to the semi-short conditions for $j = \bar{\jmath} = 0$, but there is the well known anomaly to $\mathrm{O}(g^2)$.

A remaining question is whether there are perhaps any additional shortening conditions other than those discussed here. The semi-shortening conditions which arise at unitarity bounds are clearly related to null states so that the cases considered masy be expected to be complete. As a partial test of completeness for $\mathcal{N}=2$ we have considered the decomposition of a $\mathcal{N}=4$ $\frac{1}{2}$-BPS supermultiplet in terms of $\mathcal{N}=2$ supermultiplets when we obtain

$$
\begin{aligned}
\mathcal{B}^{\frac{1}{2},\frac{1}{2}}_{[0,p,0](0,0)} \simeq\ & (p+1)\,\hat{\mathcal{B}}_{\frac{1}{2}p} \oplus \mathcal{E}_{p(0,0)} \oplus \bar{\mathcal{E}}_{-p(0,0)} \\
& \oplus (p-1)\,\hat{\mathcal{C}}_{\frac{1}{2}p-1(0,0)} \oplus p\big(\mathcal{D}_{\frac{1}{2}(p-1)(0,0)} \oplus \bar{\mathcal{D}}_{\frac{1}{2}(p-1)(0,0)}\big) \\
& \oplus \bigoplus_{k=1}^{p-2} (k+1)\big(\mathcal{B}_{\frac{1}{2}k,p-k(0,0)} \oplus \bar{\mathcal{B}}_{\frac{1}{2}k,k-p(0,0)}\big) \\
& \oplus \bigoplus_{k=0}^{p-3} (k+1)\big(\mathcal{C}_{\frac{1}{2}k,p-k-2(0,0)} \oplus \bar{\mathcal{C}}_{\frac{1}{2}k,k-p+2(0,0)}\big) \\
& \oplus \bigoplus_{k=0}^{p-4} \bigoplus_{l=0}^{p-k-4} (k+1)\,\mathcal{A}^{p}_{\frac{1}{2}k,p-k-4-2l(0,0)}\,.
\end{aligned}
\tag{7.2}
$$

All possible supermultiplets discussed in section 4 arise. We may note that, as expected, for $p=2$ the decomposition contains the $\mathcal{N}=2$ energy momentum tensor multiplet $\hat{\mathcal{C}}_{0(0,0)}$.

Acknowledgements

HO would like to thank Massimo Bianchi, Gleb Arutyunov and Emery Sokatchev for useful conversations at various stages during this work, in particular to Massimo Bianchi for emphasising the importance of understanding possible decompositions of long supermultiplets. FAD would like to thank Michael Nirschl for helpful conversations and Hendryk Pfeiffer for computer assistance in the initial stages of this work. He would also like to thank Trinity College, Cambridge and the EPSRC for financial support.

Appendix A. Two Point Functions

To construct the conformal two point functions from the results of section 2 we first note that from the definition (2.14) and taking the field $\mathcal{O}_I(t, rn)$, for $n_{\hat{a}}$ a unit vector, we may obtain

$$|\Delta\rangle_I = \mathcal{O}_I(i,0)|0\rangle\, 2^{\Delta}\,, \qquad {}_{\bar{I}}\langle\Delta| = 2^{\Delta}\langle 0|\mathcal{O}_{\bar{I}}(-i,0)\,, \tag{A.1}$$

which may be derived in the same fashion as other results obtained later. We then consider $\exp(-i\beta n_{\hat{a}}\mathcal{E}^{+}{}_{\hat{a}} - i\alpha\mathcal{E}^{+}{}_{d})|\Delta\rangle_I$. To obtain an explicit expression we first consider

$$e^{-i\alpha\mathcal{E}^{+}{}_{d}}\mathcal{O}_I(i,0)|0\rangle = \mathcal{O}_I(i\tau,0)|0\rangle f(\tau)\,. \tag{A.2}$$

From this we may derive

$$\begin{aligned}\frac{\mathrm{d}}{\mathrm{d}\alpha}\big(\mathcal{O}_I(i\tau,0)|0\rangle f(\tau)\big) =& -i[\mathcal{E}^{+}{}_{d}, \mathcal{O}_I(i\tau,0)]|0\rangle\, f(\tau)\\ =& -\tfrac{1}{2}\big((1+\tau)^2\partial_\tau + 2\Delta(1+\tau)\big)\mathcal{O}_I(i\tau,0)|0\rangle\, f(\tau)\,,\end{aligned} \tag{A.3}$$

using (2.5). By solving the differential equation we easily obtain

$$\tau + 1 = \frac{2}{1+\alpha}\,, \qquad f(\tau) = \big(\tfrac{1}{2}(\tau+1)\big)^{2\Delta}\,. \tag{A.4}$$

Similarly for

$$e^{-i\beta n_{\hat{a}}\mathcal{E}^{+}{}_{\hat{a}}}\mathcal{O}_I(i\tau_0,0)|0\rangle = \mathcal{O}_J(i\tau, rn)|0\rangle\, g^{J}{}_{I}(\tau, r)\,, \tag{A.5}$$

and the following equations are now obtained

$$\begin{aligned}\frac{\mathrm{d}}{\mathrm{d}\beta}(r - i\tau) =& -\tfrac{1}{2}\big(r - i(\tau+1)\big)^2\,,\\ \frac{\mathrm{d}}{\mathrm{d}\beta}g^{J}{}_{I}(\tau,r) =& -\Delta r\, g^{J}{}_{I}(\tau,r) + (\tau+1)(s_{0\hat{a}}n_{\hat{a}})^{J}{}_{K}g^{K}{}_{I}(\tau,r)\,.\end{aligned} \tag{A.6}$$

Hence we get

$$\begin{aligned}\exp(-i\beta n_{\hat{a}}\mathcal{E}^{+}{}_{\hat{a}} - i\alpha\mathcal{E}^{+}{}_{d})|\Delta\rangle_I = \mathcal{O}_J(i(\tau-1), rn)|0\rangle\big((\exp(\tan^{-1}\tfrac{r}{\tau}\, 2s_{0\hat{a}}n_{\hat{a}})\big)^{J}{}_{I}\big(\tfrac{1}{2}(\tau^2+r^2)\big)^{\Delta}\,,\\ r = \frac{2\beta}{\beta^2 + (1+\alpha)^2}\,, \qquad \tau = \frac{2(1+\alpha)}{\beta^2+(1+\alpha)^2}\,.\end{aligned} \tag{A.7}$$

Since ${}_{\bar{I}}\langle\Delta|\exp(-i\beta n_{\hat{a}}\mathcal{E}^{+}{}_{\hat{a}} - i\alpha\mathcal{E}^{+}{}_{d})|\Delta\rangle_I = N_{\bar{I}I}$, from (2.15) and (2.17), we then determine the two point function in the form

$$\begin{aligned}\langle 0|\mathcal{O}_{\bar{I}}(-i,0)\,\mathcal{O}_I(i(\tau-1), rn)|0\rangle = \langle 0|\mathcal{O}_{\bar{I}}(0)\,\mathcal{O}_I(x)|0\rangle = \mathcal{I}_{\bar{I}I}(x)\,\frac{1}{(x^2)^{\Delta}}\,,\\ x^a = (i\tau, rn)\,, \quad x^2 = \tau^2 + r^2\,,\end{aligned} \tag{A.8}$$

for

$$\mathcal{I}(x) = N \exp(-\tan^{-1}\tfrac{r}{\tau}\, 2 s_{0\hat{a}} n_{\hat{a}}) = \exp(\tan^{-1}\tfrac{r}{\tau}\, 2\bar{s}_{0\hat{a}} n_{\hat{a}})\, N\,, \tag{A.9}$$

using (2.18) and (2.16).

As an illustration for a four dimensional spinor field, $\mathcal{O}_I = \bar{\psi}_{\dot{\alpha}}$, $\bar{\mathcal{O}}_{\bar{I}} = \psi_\alpha$, then $s_{ab} = \frac{1}{2} i \bar{\sigma}_{[a}\sigma_{b]}$ and we may take $N = \sigma_0$ so that (A.9) gives

$$\mathcal{I}(x) = -i \frac{1}{(x^2)^{\frac{1}{2}}}\, \sigma_a x^a\,. \tag{A.10}$$

Alternatively for a vector field V_a we have $(s_{ab})^c{}_d = i(\eta_{ad}\delta^c{}_b - \eta_{bd}\delta^c{}_a)$ then using $N_{cd} = \delta_{cd}$ from (A.9) we have

$$\mathcal{I}_{cd}(x) = \eta_{cd} - 2\,\frac{x_c x_d}{x^2}\,, \tag{A.11}$$

with x_c determined from (A.8). Of course (A.11) is just the expected result given by the inversion tensor.

Appendix B. Racah Speiser Algorithm

We describe here in outline the Racah Speiser algorithm [20] which provides a succinct and simple way of decomposing tensor products[4]. A representation $R_{\underline{\Lambda}}$ with Dynkin labels $\underline{\Lambda} = [\lambda_1, \ldots, \lambda_r]$ is characterised by a highest weight vector Λ,

$$\Lambda = \sum_{i=1}^{r} \lambda_i w_i \in W \quad \Leftrightarrow \quad \lambda_i \in \mathbb{Z},\ \lambda_i \geq 0\,, \tag{B.1}$$

where w_i are the fundamental weights which are dual to the simple roots α_i, $2w_i \cdot \alpha_j / {\alpha_j}^2 = \delta_{ij}$. For a tensor product $R_{\underline{\Lambda}} \otimes R_{\underline{\Lambda}'}$ we consider the set of weights $V_{\underline{\Lambda}'} = \{\lambda\}$ for all states in the representation space for $R_{\underline{\Lambda}'}$ ($\lambda = \sum_i \lambda_i w_i$ with λ_i positive or negative integers and, allowing for multiplicity, there are $d_{\Lambda'}$ weights in $V_{\underline{\Lambda}'}$ for $d_{\Lambda'}$ the dimension of $R_{\underline{\Lambda}'}$). Then the Racah Speiser algorithm can be paraphrased by expressing the tensor product as

$$R_{\underline{\Lambda}} \otimes R_{\underline{\Lambda}'} \simeq \sum_{\lambda \in V_{\underline{\Lambda}'}} R_{\underline{\Lambda}+\underline{\lambda}} = \sum_n R_{\underline{\Lambda}_n}\,, \quad \Lambda_n \in W\,, \tag{B.2}$$

where we require

$$R_{\underline{\lambda}} = \mathrm{sign}(\sigma) R_{\underline{\lambda}^\sigma}\,, \qquad \lambda^\sigma = \sigma(\lambda + \rho) - \rho\,, \tag{B.3}$$

[4] For a more pictorial description for the case of $SU(3)$ see [33] and in a mathematical context chapter VI, 24.4, exercise 9 in [34].

for σ an element of $\mathcal{W}$, the Weyl group, the symmetry group of the root space which permutes the different possible bases of simple roots, and the Weyl vector ρ is defined by

$$\rho = \tfrac{1}{2} \sum_{\alpha \in \Phi^+} \alpha = \sum_{i=1}^{r} w_i \,, \tag{B.4}$$

for Φ^+ the set of positive roots. $\mathcal{W}$ is generated by the reflections corresponding to the simple roots

$$\sigma_i \lambda = \lambda - \frac{2\lambda \cdot \alpha_i}{{\alpha_i}^2} \, \alpha_i \,, \tag{B.5}$$

and $\mathrm{sign}(\sigma) = (-1)^{N_\sigma}$ if σ is formed from the product of N_σ elementary reflections (B.5).

For $\lambda = \sum_i \lambda_i w_i$ then $\lambda \in \partial W$ if $\lambda_i = 0$ for some i, in which case $\sigma_i \lambda = \lambda$. For any weight vector λ there is a $\sigma \in \mathcal{W}$ such that $\sigma\lambda \in W$. If $\sigma\lambda \notin \partial W$ then σ is unique, if $\sigma\lambda \in \partial W$ then there is a reflection σ_λ, with $\mathrm{sign}(\sigma_\lambda) = -$, such that $\sigma_\lambda \lambda = \lambda$. Consequently if there is a σ so that $\sigma(\lambda + \rho) \in W \setminus \partial W$ then in (B.3), for all such λ, $\lambda^\sigma \in W$ is unique. Otherwise there is a reflection σ, with $\mathrm{sign}(\sigma) = -$, such that $\lambda^\sigma = \lambda$ and in this case from (B.3)

$$R_{\underline{\lambda}} = -R_{\underline{\lambda}} = 0 \,. \tag{B.6}$$

Hence in (B.2), for any weights $\lambda \in V_{\underline{\Lambda}'}$, we may let $R_{\underline{\Lambda}+\underline{\lambda}} \to \pm R_{\underline{\Lambda}_n}$ for some unique $\Lambda_n \in W$ or alternatively $R_{\underline{\Lambda}+\underline{\lambda}} \to 0$. If $\Lambda + \lambda \in W$ then we may take, for some n, $\Lambda_n = \Lambda + \lambda$. Unless this is true for all $\lambda \in V_{\underline{\Lambda}'}$ there is a cancellation of representations in the sum over $\{\Lambda_n\}$ on the right hand side of (B.2) due to $\mathrm{sign}(\sigma) = -$ for some cases.

It is easy to check that the algorithm is compatible with dimensions. Let

$$d_{\underline{\lambda}} = \prod_{\alpha \in \Phi^+} \frac{(\lambda + \rho) \cdot \alpha}{\rho \cdot \alpha} \,. \tag{B.7}$$

For any $\Lambda \in W$ $d_{\underline{\Lambda}}$ is just the Weyl formula the dimension of the representation $R_{\underline{\Lambda}}$ with Dynkin labels $\underline{\Lambda}$. For an elementary reflection as in (B.5),

$$\sigma_i(\Phi^+ \setminus \alpha_i) = \Phi^+ \setminus \alpha_i \,, \quad \sigma_i \alpha_i = -\alpha_i \quad \Rightarrow \quad d_{\underline{\lambda}^{\sigma_i}} = -d_{\underline{\lambda}} \,, \tag{B.8}$$

since the scalar product is invariant under reflections. In general then $d_{\underline{\lambda}^\sigma} = \mathrm{sign}(\sigma) d_{\underline{\lambda}}$ and it is straightforward to see that (when (B.6) holds $d_{\underline{\lambda}} = 0$),

$$\sum_{\lambda \in V_{\underline{\Lambda}'}} d_{\underline{\Lambda}+\underline{\lambda}} = \sum_n d_{\underline{\Lambda}_n} \,. \tag{B.9}$$

For illustration for the simplest case of $SU(2)$ where, following convention, the representations are R_j, $j = 0, \frac{1}{2}, 1, \ldots$, the associated weights $V_j = \{-j, -j+1, \ldots, j\}$, and the Weyl group $\mathcal{W} = \mathbb{Z}_2$, where for $m \in V_j$, $\sigma m = -m$. In this case (B.2) gives

$$R_j \otimes R_{j'} \simeq \begin{cases} \sum_{m \in V_{j'}} R_{j+m}\,, & j' \leq j\ , \\ \sum_{m \in V_{j'}} R_{j+m} = \sum_{m \in V_j} R_{j'+m}, & j' \geq j\ , \end{cases} \tag{B.10}$$

using from (B.3),

$$R_{-n} = -R_{n-1}\,, \qquad R_{-\frac{1}{2}} = 0\,. \tag{B.11}$$

Of course this is just the standard result for tensor products in $SU(2)$.

Instead of (B.5) and (B.4), defining for $\underline{\lambda} = [\ldots, \lambda_j, \ldots]$ for $\lambda = \sum_j \lambda_j w_j$, we may write

$$\sigma_i \underline{\lambda} = [\ldots, \lambda_j - \lambda_i K_{ij}, \ldots]\,, \qquad \underline{\rho} = [1, 1, \ldots, 1]\,, \tag{B.12}$$

for $[K_{ij}]$ the Cartan matrix. For the case of $SU(4)$ then with the Cartan matrix (3.25) and with (B.3) the elementary reflections give

$$\begin{aligned} &[\lambda_1, \lambda_2, \lambda_3]^{\sigma_1} = [-\lambda_1 - 2, \lambda_1 + \lambda_2 + 1, \lambda_3]\,, \quad [\lambda_1, \lambda_2, \lambda_3]^{\sigma_3} = [\lambda_1, \lambda_2 + \lambda_3 + 1, -\lambda_3 - 2]\,, \\ &[\lambda_1, \lambda_2, \lambda_3]^{\sigma_2} = [\lambda_1 + \lambda_2 + 1, -\lambda_2 - 2, \lambda_2 + \lambda_3 + 1]\,. \end{aligned} \tag{B.13}$$

For this case $\mathcal{W} = \mathcal{S}_4$, the permutation group on four objects (in this case we have the relations $\sigma_i^2 = e$, $\sigma_1\sigma_3 = \sigma_3\sigma_1$, $\sigma_1\sigma_2\sigma_1 = \sigma_2\sigma_1\sigma_2$, $\sigma_3\sigma_2\sigma_3 = \sigma_2\sigma_3\sigma_2$). Note that

$$\begin{aligned} [\lambda_1, \lambda_2, \lambda_3]^{\sigma_1\sigma_2\sigma_1} &= [-\lambda_2 - 2, -\lambda_1 - 2, \lambda_1 + \lambda_2 + \lambda_3 + 2]\,, \\ [\lambda_1, \lambda_2, \lambda_3]^{\sigma_3\sigma_2\sigma_3} &= [\lambda_1 + \lambda_2 + \lambda_3 + 2, -\lambda_3 - 2, -\lambda_2 - 2]\,, \\ [\lambda_1, \lambda_2, \lambda_3]^{\sigma_1\sigma_2\sigma_3\sigma_2\sigma_1} &= [-\lambda_2 - \lambda_3 - 3, \lambda_2, -\lambda_1 - \lambda_2 - 3]\,. \end{aligned} \tag{B.14}$$

From (B.13) and (B.14) it is clear that we have $\underline{\lambda}^\sigma = \underline{\lambda}$ with $\mathrm{sign}\sigma = -$ if one of the following conditions apply,

$$\lambda_1 = -1\,, \quad \lambda_2 = -1\,, \quad \lambda_3 = -1\,, \quad \lambda_1 + \lambda_2 = -2\,, \quad \lambda_2 + \lambda_3 = -2\,, \quad \lambda_1 + \lambda_2 + \lambda_3 = -3\,. \tag{B.15}$$

These correspond exactly to the zeros of the dimension formula (3.29), as expected since these cases correspond to zero contributions in the tensor product expansion.

Appendix C. Tables of Representations

$\ell+2q+p+2$	$\ell+2q+p+3$
$[q,p,q]_\ell$	$[q{-}1,p,q{-}1]_{\ell+1}$,$[q{-}1,p{+}2,q{-}1]_{\ell+1}$,$4[q,p,q]_{\ell+1}$,$[q{+}1,p{-}2,q{+}1]_{\ell+1}$,$[q{+}1,p,q{+}1]_{\ell+1}$ $[q{-}1,p,q{-}1]_{\ell-1}$,$[q{-}1,p{+}2,q{-}1]_{\ell-1}$,$2[q,p,q]_{\ell-1}$,$[q{+}1,p{-}2,q{+}1]_{\ell-1}$

$\ell+2q+p+4$
$[q{-}2,p{+}2,q{-}2]_{\ell+2}$,$2[q{-}1,p,q{-}1]_{\ell+2}$,$2[q{-}1,p{+}2,q{-}1]_{\ell+2}$,$[q,p{-}2,q]_{\ell+2}$ $6[q,p,q]_{\ell+2}$,$[q,p{+}2,q]_{\ell+2}$,$2[q{+}1,p{-}2,q{+}1]_{\ell+2}$,$2[q{+}1,p,q{+}1]_{\ell+2}$,$[q{+}2,p{-}2,q{+}2]_{\ell+2}$ $[q{-}2,p,q{-}2]_\ell$,$4[q{-}2,p{+}2,q{-}2]_\ell$,$[q{-}2,p{+}4,q{-}2]_\ell$,$10[q{-}1,p,q{-}1]_\ell$,$8[q{-}1,p{+}2,q{-}1]_\ell$,$4[q,p{-}2,q]_\ell$ $15[q,p,q]_\ell$,$[q,p{+}2,q]_\ell$,$8[q{+}1,p{-}2,q{+}1]_\ell$,$2[q{+}1,p,q{+}1]_\ell$,$[q{+}2,p{-}4,q{+}2]_\ell$,$[q{+}2,p{-}2,q{+}2]_\ell$ $[q{-}2,p{+}2,q{-}2]_{\ell-2}$,$2[q{-}1,p,q{-}1]_{\ell-2}$,$[q,p{-}2,q]_{\ell-2}$,$[q,p,q]_{\ell-2}$

$\ell+2q+p+5$
$[q{-}1,p,q{-}1]_{\ell+3}$,$[q{-}1,p{+}2,q{-}1]_{\ell+3}$,$4[q,p,q]_{\ell+3}$,$[q{+}1,p{-}2,q{+}1]_{\ell+3}$,$[q{+}1,p,q{+}1]_{\ell+3}$ $[q{-}3,p{+}2,q{-}3]_{\ell+1}$,$[q{-}3,p{+}4,q{-}3]_{\ell+1}$,$2[q{-}2,p,q{-}2]_{\ell+1}$,$10[q{-}2,p{+}2,q{-}2]_{\ell+1}$,$2[q{-}2,p{+}4,q{-}2]_{\ell+1}$,$[q{-}1,p{-}2,q{-}1]_{\ell+1}$ $20[q{-}1,p,q{-}1]_{\ell+1}$,$15[q{-}1,p{+}2,q{-}1]_{\ell+1}$,$[q{-}1,p{+}4,q{-}1]_{\ell+1}$,$10[q,p{-}2,q]_{\ell+1}$,$26[q,p,q]_{\ell+1}$,$4[q,p{+}2,q]_{\ell+1}$ $[q{+}1,p{-}4,q{+}1]_{\ell+1}$,$15[q{+}1,p{-}2,q{+}1]_{\ell+1}$,$6[q{+}1,p,q{+}1]_{\ell+1}$,$2[q{+}2,p{-}4,q{+}2]_{\ell+1}$,$4[q{+}2,p{-}2,q{+}2]_{\ell+1}$,$[q{+}3,p{-}4,q{+}3]_{\ell+1}$ $[q{-}3,p{+}2,q{-}3]_{\ell-1}$,$[q{-}3,p{+}4,q{-}3]_{\ell-1}$,$2[q{-}2,p,q{-}2]_{\ell-1}$,$8[q{-}2,p{+}2,q{-}2]_{\ell-1}$ $[q{-}1,p{-}2,q{-}1]_{\ell-1}$,$15[q{-}1,p,q{-}1]_{\ell-1}$,$4[q{-}1,p{+}2,q{-}1]_{\ell-1}$,$8[q,p{-}2,q]_{\ell-1}$,$10[q,p,q]_{\ell-1}$ $[q{+}1,p{-}4,q{+}1]_{\ell-1}$,$4[q{+}1,p{-}2,q{+}1]_{\ell-1}$,$[q{+}1,p,q{+}1]_{\ell-1}$,$[q{-}1,p,q{-}1]_{\ell-3}$

$\ell+2q+p+6$
$[q,p,q]_{\ell+4}$,$[q{-}2,p,q{-}2]_{\ell+2}$,$4[q{-}2,p{+}2,q{-}2]_{\ell+2}$,$[q{-}2,p{+}4,q{-}2]_{\ell+2}$,$10[q{-}1,p,q{-}1]_{\ell+2}$,$8[q{-}1,p{+}2,q{-}1]_{\ell+2}$ $4[q,p{-}2,q]_{\ell+2}$,$15[q,p,q]_{\ell+2}$,$[q,p{+}2,q]_{\ell+2}$,$8[q{+}1,p{-}2,q{+}1]_{\ell+2}$,$2[q{+}1,p,q{+}1]_{\ell+2}$,$[q{+}2,p{-}4,q{+}2]_{\ell+2}$,$[q{+}2,p{-}2,q{+}2]_{\ell+2}$ $[q{-}4,p{+}4,q{-}4]_\ell$,$4[q{-}3,p{+}2,q{-}3]_\ell$,$2[q{-}3,p{+}4,q{-}3]_\ell$,$6[q{-}2,p,q{-}2]_\ell$,$15[q{-}2,p{+}2,q{-}2]_\ell$,$[q{-}2,p{+}4,q{-}2]_\ell$ $4[q{-}1,p{-}2,q{-}1]_\ell$,$26[q{-}1,p,q{-}1]_\ell$,$10[q{-}1,p{+}2,q{-}1]_\ell$,$[q,p{-}4,q]_\ell$,$15[q,p{-}2,q]_\ell$,$20[q,p,q]_\ell$,$[q,p{+}2,q]_\ell$ $2[q{+}1,p{-}4,q{+}1]_\ell$,$10[q{+}1,p{-}2,q{+}1]_\ell$,$2[q{+}1,p,q{+}1]_\ell$,$[q{+}2,p{-}4,q{+}2]_\ell$,$[q{+}2,p{-}2,q{+}2]_\ell$ $[q{-}2,p,q{-}2]_{\ell-2}$,$[q{-}2,p{+}2,q{-}2]_{\ell-2}$,$4[q{-}1,p,q{-}1]_{\ell-2}$,$[q,p{-}2,q]_{\ell-2}$,$[q,p,q]_{\ell-2}$

$\ell+2q+p+7$
$[q{-}1,p,q{-}1]_{\ell+3}$,$[q{-}1,p{+}2,q{-}1]_{\ell+3}$,$2[q,p,q]_{\ell+3}$,$[q{+}1,p{-}2,q{+}1]_{\ell+3}$ $[q{-}3,p{+}2,q{-}3]_{\ell+1}$,$[q{-}3,p{+}4,q{-}3]_{\ell+1}$,$2[q{-}2,p,q{-}2]_{\ell+1}$,$8[q{-}2,p{+}2,q{-}2]_{\ell+1}$,$[q{-}1,p{-}2,q{-}1]_{\ell+1}$,$15[q{-}1,p,q{-}1]_{\ell+1}$ $4[q{-}1,p{+}2,q{-}1]_{\ell+1}$,$8[q,p{-}2,q]_{\ell+1}$,$10[q,p,q]_{\ell+1}$,$[q{+}1,p{-}4,q{+}1]_{\ell+1}$,$4[q{+}1,p{-}2,q{+}1]_{\ell+1}$,$[q{+}1,p,q{+}1]_{\ell+1}$ $[q{-}3,p{+}2,q{-}3]_{\ell-1}$,$2[q{-}2,p,q{-}2]_{\ell-1}$,$2[q{-}2,p{+}2,q{-}2]_{\ell-1}$,$[q{-}1,p{-}2,q{-}1]_{\ell-1}$ $6[q{-}1,p,q{-}1]_{\ell-1}$,$[q{-}1,p{+}2,q{-}1]_{\ell-1}$,$2[q,p{-}2,q]_{\ell-1}$,$2[q,p,q]_{\ell-1}$,$[q{+}1,p{-}2,q{+}1]_{\ell-1}$

$\ell+2q+p+8$	$\ell+2q+p+9$
$[q{-}2,p{+}2,q{-}2]_{\ell+2}$,$2[q{-}1,p,q{-}1]_{\ell+2}$,$[q,p{-}2,q]_{\ell+2}$,$[q,p,q]_{\ell+2}$ $[q{-}2,p,q{-}2]_\ell$,$[q{-}2,p{+}2,q{-}2]_\ell$,$4[q{-}1,p,q{-}1]_\ell$,$[q,p{-}2,q]_\ell$,$[q,p,q]_\ell$	$[q-1,p,q-1]_{\ell+1}$

Table 4. Diagonal representations in $\mathcal{C}^{\frac{1}{4},\frac{1}{4}}_{[q,p,q](\frac{1}{2}\ell,\frac{1}{2}\ell)}$.

$p+2$	$p+3$	$p+4$
$[1,p,1]_0$	$[0,p,0]_1$,$[0,p{+}2,0]_1$ $2[1,p,1]_1$,$[2,p{-}2,2]_1$	$2[0,p,0]_2$,$[1,p{-}2,1]_2$,$[1,p,1]_2$ $[0,p,0]_0$,$[1,p{-}2,1]_0$,$2[1,p,1]_0$,$2[2,p{-}2,2]_0$,$[3,p{-}4,3]_0$

$p+5$	$p+6$	$p+7$
$[0,p,0]_3$,$[0,p{-}2,0]_1$,$2[0,p,0]_1$,$4[1,p{-}2,1]_1$ $[1,p,1]_1$,$[2,p{-}4,2]_1$,$[2,p{-}2,2]_1$	$[1,p{-}2,1]_2$,$2[0,p{-}2,0]_0$ $[1,p{-}4,1]_0$,$[1,p{-}2,1]_0$	$[0,p-2,0]_1$

Table 5. Diagonal representations in $\mathcal{B}^{\frac{1}{4},\frac{1}{4}}_{[1,p,1](0,0)}$.

$\ell+p+4$	$\ell+p+5$
$[1,p,1]_\ell$	$[0,p,0]_{\ell+1}$,$[0,p{+}2,0]_{\ell+1}$,$4[1,p,1]_{\ell+1}$,$[2,p{-}2,2]_{\ell+1}$,$[2,p,2]_{\ell+1}$ $[0,p,0]_{\ell-1}$,$[0,p{+}2,0]_{\ell-1}$,$2[1,p,1]_{\ell-1}$,$[2,p{-}2,2]_{\ell-1}$

$\ell+p+6$
$2[0,p,0]_{\ell+2}$,$2[0,p{+}2,0]_{\ell+2}$,$[1,p{-}2,1]_{\ell+2}$,$6[1,p,1]_{\ell+2}$,$[1,p{+}2,1]_{\ell+2}$,$2[2,p{-}2,2]_{\ell+2}$,$2[2,p,2]_{\ell+2}$,$[3,p{-}2,3]_{\ell+2}$ $6[0,p,0]_\ell$,$4[0,p{+}2,0]_\ell$,$4[1,p{-}2,1]_\ell$,$13[1,p,1]_\ell$,$[1,p{+}2,1]_\ell$,$8[2,p{-}2,2]_\ell$,$2[2,p,2]_\ell$,$[3,p{-}4,3]_\ell$,$[3,p{-}2,3]_\ell$ $2[0,p,0]_{\ell-2}$,$[1,p{-}2,1]_{\ell-2}$,$[1,p,1]_{\ell-2}$,

$\ell+p+7$
$[0,p,0]_{\ell+3}$,$[0,p{+}2,0]_{\ell+3}$,$4[1,p,1]_{\ell+3}$,$[2,p{-}2,2]_{\ell+3}$,$[2,p,2]_{\ell+3}$ $[0,p{-}2,0]_{\ell+1}$,$11[0,p,0]_{\ell+1}$,$8[0,p{+}2,0]_{\ell+1}$,$[0,p{+}4,0]_{\ell+1}$,$10[1,p{-}2,1]_{\ell+1}$,$22[1,p,1]_{\ell+1}$,$4[1,p{+}2,1]_{\ell+1}$ $[2,p{-}4,2]_{\ell+1}$,$15[2,p{-}2,2]_{\ell+1}$,$6[2,p,2]_{\ell+1}$,$2[3,p{-}4,3]_{\ell+1}$,$4[3,p{-}2,3]_{\ell+1}$,$[4,p{-}4,4]_{\ell+1}$ $[0,p{-}2,0]_{\ell-1}$,$8[0,p,0]_{\ell-1}$,$[0,p{+}2,0]_{\ell-1}$,$8[1,p{-}2,1]_{\ell-1}$,$8[1,p,1]_{\ell-1}$,$[2,p{-}4,2]_{\ell-1}$,$4[2,p{-}2,2]_{\ell-1}$,$[2,p,2]_{\ell-1}$,$[0,p,0]_{\ell-3}$

$\ell+p+8$
$[1,p,1]_{\ell+4}$,$6[0,p,0]_{\ell+2}$,$4[0,p{+}2,0]_{\ell+2}$,$4[1,p{-}2,1]_{\ell+2}$,$13[1,p,1]_{\ell+2}$,$[1,p{+}2,1]_{\ell+2}$ $8[2,p{-}2,2]_{\ell+2}$,$2[2,p,2]_{\ell+2}$,$[3,p{-}4,3]_{\ell+2}$,$[3,p{-}2,3]_{\ell+2}$ $4[0,p{-}2,0]_\ell$,$12[0,p,0]_\ell$,$4[0,p{+}2,0]_\ell$,$[1,p{-}4,1]_\ell$,$15[1,p{-}2,1]_\ell$,$15[1,p,1]_\ell$,$[1,p{+}2,1]_\ell$ $2[2,p{-}4,2]_\ell$,$10[2,p{-}2,2]_\ell$,$2[2,p,2]_\ell$,$[3,p{-}4,3]_\ell$,$[3,p{-}2,3]_\ell$,$2[0,p,0]_{\ell-2}$,$[1,p{-}2,1]_{\ell-2}$,$[1,p,1]_{\ell-2}$

$\ell+p+9$
$[0,p,0]_{\ell+3}$,$[0,p{+}2,0]_{\ell+3}$,$2[1,p,1]_{\ell+3}$,$[2,p{-}2,2]_{\ell+3}$ $[0,p{-}2,0]_{\ell+1}$,$8[0,p,0]_{\ell+1}$,$[0,p{+}2,0]_{\ell+1}$,$8[1,p{-}2,1]_{\ell+1}$,$8[1,p,1]_{\ell+1}$,$[2,p{-}4,2]_{\ell+1}$,$4[2,p{-}2,2]_{\ell+1}$,$[2,p,2]_{\ell+1}$ $[0,p{-}2,0]_{\ell-1}$,$3[0,p,0]_{\ell-1}$,$[0,p{+}2,0]_{\ell-1}$,$2[1,p{-}2,1]_{\ell-1}$,$2[1,p,1]_{\ell-1}$,$[2,p{-}2,2]_{\ell-1}$

$\ell+p+10$	$\ell+p+11$
$2[0,p,0]_{\ell+2}$,$[1,p{-}2,1]_{\ell+2}$,$[1,p,1]_{\ell+2}$ $2[0,p,0]_\ell$,$[1,p{-}2,1]_\ell$,$[1,p,1]_\ell$	$[0,p,0]_{\ell+1}$

Table 6. Diagonal representations in $\mathcal{C}^{\frac{1}{4},\frac{1}{4}}_{[1,p,1](\frac{1}{2}\ell,\frac{1}{2}\ell)}$.

$\ell+4$	$\ell+5$	$\ell+6$	$\ell+7$
$[0,2,0]_\ell$	$2[0,2,0]_{\ell+1}$,$[1,0,1]_{\ell+1}$ $[1,2,1]_{\ell+1}$,$[1,0,1]_{\ell-1}$	$[0,0,0]_{\ell+2}$,$3[0,2,0]_{\ell+2}$,$[0,4,0]_{\ell+2}$ $2[1,0,1]_{\ell+2}$,$2[1,2,1]_{\ell+2}$,$[2,0,2]_{\ell+2}$ $[0,0,0]_\ell$,$[0,2,0]_\ell$,$3[1,0,1]_\ell$,$[2,0,2]_\ell$ $[0,0,0]_{\ell-2}$	$2[0,2,0]_{\ell+3}$,$[1,0,1]_{\ell+3}$,$[1,2,1]_{\ell+3}$ $[0,0,0]_{\ell+1}$,$2[0,2,0]_{\ell+1}$,$4[1,0,1]_{\ell+1}$ $[1,2,1]_{\ell+1}$,$[2,0,2]_{\ell+1}$ $[0,0,0]_{\ell-1}$,$[1,0,1]_{\ell-1}$

$\ell+8$	$\ell+9$	$\ell+10$
$[0,2,0]_{\ell+4}$,$[0,0,0]_{\ell+2}$,$[0,2,0]_{\ell+2}$,$3[1,0,1]_{\ell+2}$,$[2,0,2]_{\ell+2}$ $[0,0,0]_\ell$,$[0,2,0]_\ell$,$[1,0,1]_\ell$	$[1,0,1]_{\ell+3}$ $[0,0,0]_{\ell+1}$,$[1,0,1]_{\ell+1}$	$[0,0,0]_{\ell+2}$

Table 7. Diagonal representations in $\mathcal{C}^{\frac{1}{2},\frac{1}{2}}_{[0,2,0](\frac{1}{2}\ell,\frac{1}{2}\ell)}$.

$2q+p+2$	$2q+p+3$
$[q,p,q]_0$	$[q-1,p,q-1]_1$,$[q-1,p+2,q-1]_1$,$4[q,p,q]_1$,$[q+1,p-2,q+1]_1$,$[q+1,p,q+1]_1$

$2q+p+4$
$[q-2,p+2,q-2]_2$,$2[q-1,p,q-1]_2$,$2[q-1,p+2,q-1]_2$,$[q,p-2,q]_2$ $6[q,p,q]_2$,$[q,p+2,q]_2$,$2[q+1,p-2,q+1]_2$,$2[q+1,p,q+1]_2$,$[q+2,p-2,q+2]_2$ $[q-2,p,q-2]_0$,$[q-2,p+2,q-2]_0$,$[q-2,p+4,q-2]_0$,$4[q-1,p,q-1]_0$,$4[q-1,p+2,q-1]_0$,$[q,p-2,q]_0$ $8[q,p,q]_0$,$[q,p+2,q]_0$,$4[q+1,p-2,q+1]_0$,$2[q+1,p,q+1]_0$,$[q+2,p-4,q+2]_0$,$[q+2,p-2,q+2]_0$

$2q+p+5$
$[q-1,p,q-1]_3$,$[q-1,p+2,q-1]_3$,$4[q,p,q]_3$,$[q+1,p-2,q+1]_3$,$[q+1,p,q+1]_3$ $[q-3,p+2,q-3]_1$,$[q-3,p+4,q-3]_1$,$2[q-2,p,q-2]_1$,$8[q-2,p+2,q-2]_1$,$2[q-2,p+4,q-2]_1$,$[q-1,p-2,q-1]_1$ $15[q-1,p,q-1]_1$,$13[q-1,p+2,q-1]_1$,$[q-1,p+4,q-1]_1$,$8[q,p-2,q]_1$,$22[q,p,q]_1$,$4[q,p+2,q]_1$ $[q+1,p-4,q+1]_1$,$13[q+1,p-2,q+1]_1$,$6[q+1,p,q+1]_1$,$2[q+2,p-4,q+2]_1$,$4[q+2,p-2,q+2]_1$,$[q+3,p-4,q+3]_1$

$2q+p+6$
$[q,p,q]_4$,$[q-2,p,q-2]_2$,$4[q-2,p+2,q-2]_2$,$[q-2,p+4,q-2]_2$,$10[q-1,p,q-1]_2$,$8[q-1,p+2,q-1]_2$ $4[q,p-2,q]_2$,$15[q,p,q]_2$,$[q,p+2,q]_2$,$8[q+1,p-2,q+1]_2$,$2[q+1,p,q+1]_2$,$[q+2,p-4,q+2]_2$,$[q+2,p-2,q+2]_2$ $[q-4,p+4,q-4]_0$,$2[q-3,p+2,q-3]_0$,$2[q-3,p+4,q-3]_0$,$3[q-2,p,q-2]_0$,$8[q-2,p+2,q-2]_0$,$[q-2,p+4,q-2]_0$ $2[q-1,p-2,q-1]_0$,$12[q-1,p,q-1]_0$,$6[q-1,p+2,q-1]_0$,$[q,p-4,q]_0$,$8[q,p-2,q]_0$,$11[q,p,q]_0$,$[q,p+2,q]_0$ $2[q+1,p-4,q+1]_0$,$6[q+1,p-2,q+1]_0$,$2[q+1,p,q+1]_0$,$[q+2,p-4,q+2]_0$,$[q+2,p-2,q+2]_0$

$2q+p+7$
$[q-1,p,q-1]_3$,$[q-1,p+2,q-1]_3$,$2[q,p,q]_3$,$[q+1,p-2,q+1]_3$ $[q-3,p+2,q-3]_1$,$[q-3,p+4,q-3]_1$,$2[q-2,p,q-2]_1$,$8[q-2,p+2,q-2]_1$,$[q-1,p-2,q-1]_1$,$15[q-1,p,q-1]_1$ $4[q-1,p+2,q-1]_1$,$8[q,p-2,q]_1$,$10[q,p,q]_1$,$[q+1,p-4,q+1]_\ell$,$4[q+1,p-2,q+1]_1$,$[q+1,p,q+1]_1$

$2q+p+8$	$2q+p+9$
$[q-2,p+2,q-2]_2$,$2[q-1,p,q-1]_2$,$[q,p-2,q]_2$,$[q,p,q]_2$ $[q-2,p,q-2]_0$,$[q-2,p+2,q-2]_0$,$4[q-1,p,q-1]_0$,$[q,p-2,q]_0$,$[q,p,q]_0$	$[q-1,p,q-1]_1$

Table 8. Diagonal representations in $\mathcal{C}^{\frac{1}{4},\frac{1}{4}}_{[q,p,q](0,0)}$.

$p+2$	$p+3$	$p+4$
$[0,p,0]_0$	$2[0,p,0]_1$,$[1,p-2,1]_1$ $[1,p,1]_1$	$[0,p-2,0]_2$,$3[0,p,0]_2$,$[0,p+2,0]_2$,$2[1,p-2,1]_2$,$2[1,p,1]_2$,$[2,p-2,2]_2$ $[0,p,0]_0$,$2[1,p-2,1]_0$,$[2,p-4,2]_0$,$[2,p-2,2]_0$

$p+5$	$p+6$
$2[0,p,0]_3$,$[1,p-2,1]_3$,$[1,p,1]_3$ $2[0,p-2,0]_1$,$2[0,p,0]_1$,$[1,p-4,1]_1$,$6[1,p-2,1]_1$ $[1,p,1]_1$,$2[2,p-4,2]_1$,$2[2,p-2,2]_1$,$[3,p-4,3]_1$	$[0,p,0]_4$,$[0,p-2,0]_2$,$[0,p,0]_2$,$4[1,p-2,1]_2$,$[2,p-4,2]_2$,$[2,p-2,2]_2$ $[0,p-4,0]_0$,$3[0,p-2,0]_0$,$[0,p,0]_0$ $2[1,p-4,1]_0$,$2[1,p-2,1]_0$,$[2,p-4,2]_0$

$p+7$	$p+8$
$[1,p-2,1]_3$,$2[0,p-2,0]_1$,$[1,p-4,1]_1$,$[1,p-2,1]_1$	$[0,p-2,0]_2$

Table 9. Diagonal representations in $\mathcal{C}^{\frac{1}{2},\frac{1}{2}}_{[0,p,0](0,0)}$.

References

[1] E. D'Hoker, S.D. Mathur, A. Matsusis and L. Rastelli, *The Operator Product Expansion of $N = 4$ SYM and the 4-point Functions of Supergravity*, Nucl. Phys. B589 (2000) 38, hep-th/9911222.

[2] G. Arutyunov, S. Frolov and A.C. Petkou, *Operator Product Expansion of the Lowest Weight CPOs in $\mathcal{N} = 4$ SYM_4 at Strong Coupling*, Nucl. Phys. B586 (2000) 547, hep-th/0005182; (E) Nucl. Phys. B609 (2001) 539.

[3] G. Arutyunov, S. Frolov and A.C. Petkou, *Perturbative and instanton corrections to the OPE of CPOs in $\mathcal{N} = 4$ SYM_4*, Nucl. Phys. B602 (2001) 238, hep-th/0010137; (E) Nucl. Phys. B609 (2001) 540.

[4] L. Hoffmann, L. Mesref and W. Rühl, *Conformal partial wave analysis of AdS amplitudes for dilaton-axion four-point functions*, Nucl. Phys. B608 (2001) 177, hep-th/0012153.

[5] G. Arutyunov, B. Eden, A.C. Petkou and E. Sokatchev, *Exceptional non-renormalization properties and OPE analysis of chiral four-point functions in $\mathcal{N} = 4$ SYM_4*, hep-th/0103230.

[6] B. Eden and E. Sokatchev, *On the OPE of 1/2 BPS Short Operators in $N = 4$ $SCFT_4$*, Nucl. Phys. B618 (2001) 259, hep-th/0106249.

[7] F.A. Dolan and H. Osborn, *Superconformal symmetry, correlation functions and the operator product expansion*, Nucl. Phys. B629 (2002) 3, hep-th/0112251.

[8] V.K. Dobrev and V.B. Petkova, *All positive energy unitary irreducible representations of extended conformal symmetry*, Phys. Lett. 162B (1985) 127;
V.K. Dobrev and V.B. Petkova, *Group-theoretical approach to extended conformal supersymmetry: function space realizations and invariant differential operators*, Fortschr. der Phys. 35 (1987) 537.

[9] S. Minwalla, *Restrictions imposed by superconformal invariance on quantum field theories*, Adv. Theor. Math. Phys. 2 (1998) 781, hep-th/9712074.

[10] S. Ferrara and A. Zaffaroni, *Superconformal Field Theories, Multiplet Shortening, and the $AdS_5/SCFT_4$ Correspondence*, Proceedings of the Conférence Moshé Flato 1999, vol. 1, ed. G. Dito and D. Sternheimer, Kluwer Academic Publishers (2000), hep-th/9908163.

[11] M. Bianchi, S. Kovacs, G. Rossi and Y.S. Stanev, *Properties of the Konishi multiplet in $\mathcal{N} = 4$ SYM theory*, JHEP 0105 (2001) 042, hep-th/0104016.

[12] S. Penati and A. Santambrogio, *Superspace approach to anomalous dimensions in $\mathcal{N} = 4$ SYM*, Nucl. Phys. B614 (2001) 367, hep-th/0107071.

[13] M. Bianchi, B. Eden, G. Rossi and Y.S. Stanev, *On operator mixing in $\mathcal{N} = 4$ SYM*, hep-th/0205321.

[14] G. Arutyunov, B. Eden and E. Sokatchev, *On non-renormalization and OPE in Superconformal Field Theories*, Nucl. Phys. B619 (2001) 359, hep-th/0105254.

[15] M. Günaydin and N. Marcus, *The spectrum of the S^5 compactification of the chiral $N = 2$, $D = 10$ supergravity and the unitary supermultiplets of $U(2, 2/4)$*, Class. and Quantum Gravity, 2 (1985) L11.

[16] M. Günaydin, D. Minic and M. Zagermann, *4D doubleton conformal theories, CPT and IIB strings on $AdS_5 \times S_5$*, Nucl. Phys. B534 (1998) 96, (E) B538 (1999), 531, hep-th/9806042;
M. Günaydin, D. Minic and M. Zagermann, *Novel supermultiplets of $SU(2, 2|4)$ and the AdS_4/CFT_4 duality*, Nucl. Phys. B544 (1999) 737, hep-th/9810226.

[17] S. Ferrara and E. Sokatchev, *Short representations of $SU(2, 2|N)$ and harmonic superspace analyticity*, Lett. Math. Phys. 52 (2000) 247, hep-th/9912168;
L. Andrianopoli, S. Ferrara, E. Sokatchev and B. Zupnik, *Shortening of primary operators in N-extended $SCFT_4$ and harmonic superspace analyticity*, Adv. Theor. Math. Phys. 3 (1999) 1149, hep-th/9912007.

[18] S. Ferrara and E. Sokatchev, *Superconformal interpretation of BPS states in AdS geometries*, Int. J. Theor. Phys. 40 (2001) 935, hep-th/0005151.

[19] P.J. Heslop and P.S. Howe, *On harmonic superspaces and superconformal fields in four-dimensions*, Class. Quant. Grav. 17 (2000) 3743, hep-th/0005135.

[20] J. Fuchs and C Schweigert, *Symmetries, Lie Algebras and Representations*, Cambridge University Press, Cambridge, 1997.

[21] M.F. Sohnius, *The multiplet of currents for $N = 2$ extended supersymmetry*, Phys. Lett. 81B (1979) 8.

[22] K. Dasgupta, M.M. Sheikh-Jabbari and M. Van Raamsdonk, *Protected multiplets of M-theory on a plane wave*, hep-th/0207050.

[23] N. Kim aand J Plefka, *On the spectrum of PP-wave matrix theory*, hep-th/0207034;
N. Kim and J-H. Park, *Superalgebras for M-theory on a pp-wave*, hep-th/0207061.

[24] D. Anselmi, *Theory of higher spin tensor currents and central charges*, Nucl. Phys. B541 (1999) 323, hep-th/9808004.

[25] S.E. Konstein, M.A. Vasiliev and V.N. Zaikin, *Conformal higher spin currents in any dimension and AdS/CFT correspondence*, JHEP 0012 (2000) 018, hep-th/0010239.

[26] K. Intriligator, *Bonus Symmetries of $\mathcal{N} = 4$ Super-Yang-Mills Correlation Functions via AdS Duality*, Nucl. Phys. B551 (1999) 575, hep-th/9811047.

[27] K. Intriligator and W. Skiba, *Bonus Symmetry and the Operator Product Expansion of $\mathcal{N} = 4$ Super-Yang-Mills*, Nucl. Phys. B559 (1999) 165, hep-th/9905020.

[28] M. Nirschl and H. Osborn, in preparation, H. Osborn, talk at the 35th International Symposium Ahrenshoop, August 2002.

[29] B. Eden, P.S. Howe, C. Schubert, E. Sokatchev and P.C. West, *Extremal correlators in four-dimensional SCFT*, Phys. Lett. B472 (2000) 323, hep-th/9910150;
J. Erdmenger and M. Pérez-Victoria, *Non-renormalization of next-to-extremal correlators in $\mathcal{N} = 4$ SYM and the AdS/CFT correspondence*. Phys. Rev. D62 (2000) 045008, hep-th/9912250;
E. D'Hoker, J. Erdmenger, D.Z. Freedman and M. Pérez-Victoria, *Near-extremal correlators and vanishing supergravity couplings in AdS/CFT*, Nucl. Phys. B589 (2000) 3, hep-th/0003218;
B. Eden, P.S. Howe, E. Sokatchev and P.C. West, *Extremal and next-to-extremal n-point correlators in four dimensional SCFT*, Phys. Lett. B494 (2000) 141, hep-th/0004102.

[30] A.V. Ryzhov, *Quarter BPS operators in $\mathcal{N} = 4$ SYM*, JHEP 0111 (2001) 046, hep-th/0109064;
E. D'Hoker and A.V. Ryzhov, *Three point functions of quarter BPS operators in $\mathcal{N} = 4$ SYM*, JHEP 0202 (2002) 047, hep-th/0109065.

[31] P.J. Heslop and P.S. Howe, *A note on composite operators in $N = 4$ SYM*, Phys. Lett. 516B (2001) 367, hep-th/0106238.

[32] G. Arutyunov, S. Penati, A.C. Petkou, A. Santambrogio and E. Sokatchev, *Non protected operators in $\mathcal{N} = 4$ SYM and multiparticle states of AdS_5 SUGRA*, hep-th/0206020.

[33] J.J. de Swart, *The Octet Model and its Clebsch-Gordan Coefficients*, Rev. Mod. Phys. 35 (1964) 916.

[34] J.E. Humphreys, *Introduction to Lie Algebras and Representation Theory*, Springer-Verlag, New York, third revised printing, 1980.